

Supplementa to Issue 7-8/2015

SCS Fall Meeting 2015

Poster Abstracts

Session of Medicinal Chemistry & Chemical Biology

September 4, 2015
Ecole Polytechnique Fédérale de Lausanne (EPFL)
<http://scg.ch/fallmeeting/2015>

Siglec-8 - A Novel Target For AsthmaB. S. Kroezen¹, S. Rabbani¹, B. Ernst^{1*}¹University of Basel

Sialic acid-binding immunoglobulin-like lectins (siglecs) are members of the immunoglobulin family. They are predominantly found on cells of the immune system. Up to now, 15 siglecs have been identified.¹ Among them is Siglec-8, which is selectively expressed on human eosinophils and regulates their survival.²⁻⁴ Crosslinking Siglec-8 with specific glycan ligands or antibodies induces apoptosis.⁵ In some diseases, such as asthma or chronic rhinosinusitis, an excessive amount of eosinophils is produced, causing an inflammatory reaction. Thus, targeting Siglec-8 provides a unique opportunity to control such allergic reactions.

Using a glycan array, the tetrasaccharide 6'-sulfo-sLe^x was identified as a ligand of Siglec-8.⁶ However, it exhibits poor drug-like properties and its synthesis is laborious and of high complexity. Therefore, we synthesized a number of mono- and oligosaccharides derived from the scaffold of 6'-sulfo-sLe^x and determined their affinities for Siglec-8.

Herewith, the potent disaccharide Neu5Ac α 2-3(6-O-sulfo)Gal was identified, displaying a relative IC₅₀ of 2.2 with respect to the natural 6'-sulfo-sLe^x. Moreover, the affinity of the disaccharide ligand was significantly reduced when small modifications to this structure were made. For example, when the O-glycosidic bond was changed from α 2-3 to a α 2-6 linkage, the sulfate-group was removed or only the corresponding monosaccharide fragments were evaluated, almost no binding to Siglec-8 was observed.

These results suggest that the scaffold of Neu5Ac α 2-3(6-O-sulfo)Gal contains the essential epitope of 6'-sulfo-sLe^x required for Siglec-8 binding. Based on this information, a new family of Siglec-8 ligands is currently synthesized.

[1] Paul R. Crocker, James C. Paulson, Ajit Varki, *Nature Rev. immunol.*, **2007**, 7, 255-266.

[2] Takumi Kiwamoto, Norihito Kawasaki, James C. Paulson, Bruce S. Bochner, *Pharmacol. Therapeut.*, **2012**, 135, 327-336.

[3] Helen Floyd, *et al.*, *J. Biol. Chem.*, **2000**, 275, 861-866.

[4] Kristine K. Kikly, *et al.*, *J. Biol. Chem.*, **2000**, 105, 1093-1100.

[5] Esra Nutku, *et al.*, *Blood*, **2003**, 101, 5014-5020.

[6] Bruce S. Bochner, *et al.*, *J. Biol. Chem.*, **2005**, 280, 4307-4312.

Chemically defined chromatin and protein engineering via EPL to study histone ubiquitination on the single molecule level

A.L. Bachmann¹

¹EPF Lausanne

The epigenetic landscape, essentially defined by histone post-translational modification marks (PTMs) and different DNA methylation states, is crucial for proper packaging of DNA into eukaryotic cells and dynamic regulation of DNA during transcription, DNA damage repair (DDR) and DNA replication. Many histone PTMs have been associated with an accessible, decondensed chromatin state (acetylation, H2B ubiquitination)^[1] or a rather dense state of chromatin and chromatin silencing (H3K9/27 methylation). In the past, proteins that interact with PTMs have been studied with biochemical methods providing low dynamic resolution and suffering from ensemble averaging.

Here, we aim to reveal the dynamic behavior of effector protein interactions with chromatin fibers of defined modification states on the single-molecule level, using total internal fluorescence microscopy (TIRFM). Our laboratory has extensive expertise in semi-synthesis of chromatin arrays containing a defined set of PTMs. Via expressed protein ligation (EPL)^[2] of recombinantly expressed histones and PTMs on a synthetic histone tail we achieve traceless modification of histones, as H3K9me3 or H2BK120ub^[1]. We combine this approach with an EPL based labelling strategy to attach ultra-bright fluorophores to expressed proteins. Thus, we are able to perform single-molecule measurements (*in vitro*) using a precisely defined system of labelled proteins and chromatin arrays. In particular, we used this system to examine different binding models of HP1 α protein to chromatin.

We aim to extend our method to study enzymatic protein-chromatin interactions by focusing on histone ubiquitination. We fluorescently label a key E3 ubiquitin ligase in the the DDR, RNF168, to directly measure its dynamic association with the repressed or active chromatin state with different PTM signatures, on the single-molecule level. In parallel we will determine ubiquitination kinetics in a separated emission channel using photoactivatable fluorophores. This will put us in the position to measure chromatin state dependent ubiquitination events and determine the contribution of other PTMs/ubiquitin marks towards ubiquitin installation on chromatin.

[1] Fierz, B., et al., Nat Chem Biol, **2011**, 7(2), 113-119

[2] Vila-Perello M., et al., J Am Chem Soc, **2013**, 135(1), 286-292

Engineering of a specific probe for the visualization and analysis of bivalent epigenetic marks in living cellsA. Delachat¹, H. Pick¹, N. Guidotti¹, B. Fierz^{1*}¹EPF Lausanne

In embryonic stem (ES) cells, promoters of developmentally important genes are marked with the combination of repressive (H3 lysine 27 methylation, H3K27me3) and activating (H3 lysine 4 methylation, H3K4me3) post-translational modifications (PTMs) of histones, called bivalent marks. Bivalent chromatin domains are believed to play an important role in maintaining the pluripotency of ES cells but have been observed in cancer cells as well. Bivalent domains have thus far mostly been studied by methods involving fixed, non-living cells or requiring lysis of large number of cells and by employing specific antibodies.

Here, we combine chemical chromatin synthesis (using expressed protein ligation) and protein design to develop a genetically encoded probe that binds specifically to bivalent nucleosomes. This probe is designed to weakly interact with singly modified nucleosomes but it strongly binds to nucleosomes having both H3K4me3 and H3K27me3 marks. This sensor platform allows us to image co-existing H3K4me3 and H3K27me3 at the level of single live cells, to follow bivalent chromatin through the cell-cycle and differentiation, and to re-write the histone code.

We thus designed, expressed and optimized a series of genetically encoded multivalent protein probes consisting of a sensor (fluorescent protein), inserted between two histone PTM binding domains which cooperatively detect the bivalent marks. The binding properties of these constructs were assessed *in vitro* for different trimethyl-marks, using a library of chemically synthesized nucleosomes carrying defined combinations of H3K4me3 and H3K27me3.


Expression and subsequent observation by fluorescence confocal microscopy of functional constructs allowed the localization of bivalent chromatin in stem cells and cancer cells. These tools now allow us to directly monitor the dynamics of bivalent PTMs in living cells.

The Power of the 'SCS': Improving the Pharmacological Properties of Peptide Therapeutics

C. Kourra¹, N. Cramer^{1*}

¹EPF Lausanne

Cyclic peptide therapeutics (1) fill the gap between small molecules (5000 Da) as a novel class of drugs, combining the advantages of both in terms of high selectivity, bioavailability, synthetic accessibility and low toxicity.[1] However, their fast clearance times pose a serious limitation to their utility.[2] Several strategies exist to stabilize these peptides through local or global constraints, side chain functionalisation or amide bond surrogates.[3],[4]



The replacement of easily reducible disulfide bonds, which are commonly found in several peptide structures, by adding a linker moiety,[5] would allow us to obtain non-reducible and supposedly more stable analogues (2). We aim to maintain or even improve the pharmacological properties of known peptide drugs without negatively affecting their receptor selectivity and activity.

[1] M. Gongora-Benitez, J. Tulla-Puche, F. Albericio, *Chem. Rev.* **2014**, 114, 901-926;

[2] B. Arunachalam, U. T. Phan, H. J. Geuze, P. Cresswell, *Proc. Natl. Acad. Sci. (PNAS)* **2000**, 97, 745-750

[3] Y. H. Lau, P. Andrade, Y. Wu, D. R. Spring, *Chem. Soc. Rev.* **2015**, 44, 91-102

[4] A. K. Yudin, *Chem. Sci.* **2015**, 6, 30-49

[5] W. F. DeGrado, D. C. Greenbaum *et al.*, *J. Am. Chem. Soc.* **2012**, 134, 17704-17713

An “in vivo” temperature dependence study of the protoporphyrine IX delayed fluorescence lifetime while measuring the oxygen partial pressure.

E. L. Gerelli¹, V. Huntosova¹, J. Joniova¹, G. Wagnières^{1*}

¹EPF Lausanne

Oxygen partial pressure (pO₂) in biological tissues is central in the destruction mechanisms at play during radiotherapy and photodynamic therapy (PDT). Thus, monitoring this parameter in real time can be useful to predict the outcome of these therapeutic approaches.

The triplet state of most porphyrin -photosensitizers (PSs) used in PDT is quenched by molecular oxygen. Consequently, measurement of this state's lifetime yields information on the surrounding pO₂ via the Stern–Volmer relationship [1]. Unfortunately, the very low brightness of these PSs' phosphorescence lifetime makes it difficult to use it in clinical settings. Our original approach is to measure the lifetime of the delayed fluorescence (DF) resulting from inverse intersystem crossing (iISC) transition taking place between the triplet and singlet states of the PS. By discriminating the prompt fluorescence (PF) and the -DF of an endogenous PS with an optical fiber-based time-gated system, Piffaretti and *al.*[1] reported the first real-time *in vivo* monitoring of the radiative dose in PDT. This study was based on the measurement of the protoporphyrin IX (PPIX) DF lifetime, a PS in extensive clinical use. Our team also reported a correlation between pO₂ depletion and vascular damages induced by PDT[3]. However, temperature is known to have a significant effect on the Stern-Volmer constants in most situations. This is likely to be important in the case of PPIX's DF since the iISC is a thermal process[4]. Consequently, the temperature dependence of DF lifetime may contribute to the fluctuations reported by Piffaretti and *al.*[3] and must be known to clearly define the parameters linking PPIX's DF lifetime to vascular damages. We investigated the temperature influence on PPIX's DF lifetime *in vivo* and *in vitro*.

The *in vivo* study was carried out by measuring PPIX's DF lifetime after topical application of amino-levulinic acid, a PPIX precursor, on the chorio-allantoic membrane of fertilized chick's eggs at various physiological temperatures. Our results give insights on improvements of PDT dosimetry, in particular when the treated organ is subject to relatively important temperature fluctuations, such as cutaneous tissue.

[1] F. Piffaretti and *al.*, *J. Biomedical Optics.*, **2011**, 16, 037005.

[2] M. Sinaasappel and C. Ince, *J. Appl. Physiol.*, **1996**, 81, 2297–303.

[3] F. Piffaretti and *al.*, *J. Biomedical Optics.*, **2012**, 17, 115007.

[4] M. Scholz and *al.*, *Photochem. Photobiol.*, **2013**, 12, 1873–84.

HP1 α dynamic binding to different compaction states of chromatinL. Bryan¹, B. Fierz^{1*}¹EPF Lausanne

The genetic material of eukaryotic cells is tightly packed into the nucleus in the form of chromatin. The building blocks of chromatin are nucleosomes, composed of 145-147 basepairs of DNA wrapped around eight histone proteins, which display post-translational modifications (PTMs) on their tails, such as methylation, acetylation, phosphorylation or ubiquitination. Effectors or reader proteins are known to dynamically interact with these histone modifications. One example is the heterochromatin protein 1 (HP1) family, which binds to a specific PTM, histone H3 trimethylated at lysine 9 (H3K9me3), in a multivalent fashion and is involved in chromatin compaction, gene repression, telomere maintenance and DNA repair.

A single-molecule method using total internal reflection microscopy was recently developed in our lab to measure HP1 α dynamics with chemically modified chromatin arrays, carrying defined levels of H3K9me3¹. From these measurements, binding to individual chromatin arrays was observed to be very variable, possibly reflecting the different compaction states of the chromatin fibers.

In order to reveal the effect of fiber conformation on multivalent chromatin binding, we probe the binding dynamics of HP1 α to chromatin in various compaction states. We thus engineered H3K9me3 containing chromatin in a condensed state, through the addition of histone H1, and in a decondensed state, by inserting histone H4 containing an acetylated lysine 16.

HP1 α interaction studies on the single-molecule scale currently allow us to directly probe the effects of chromatin fiber compaction on effector binding dynamics. Taken together, these results will lead to a greater understanding of molecular mechanisms involved in epigenetic gene repression.


[1] Sinan Kilic, Andreas Bachmann, Louise Bryan, Beat Fierz, *Nature communications*, (in press)

Tackling Malaria by Inhibiting the SHMT Enzyme

G. Schwartz¹, M. Witschel², M. Rottmann³, P. Chaiyen⁴, U. Leartsakulpanich⁴, K. White⁵, F. Diederich^{1*}

¹ETH Zurich, ²BASF AG, ³Swiss Tropical and Public Health Institute, Basel, ⁴National Center for Genetic Engineering and Biotechnology, Thailand, ⁵Monash Institute of Pharmaceutical Sciences

Malaria is mainly caused by the parasite *Plasmodium falciparum*. Due to the emergence of drug-resistant strains, there is an urgent need of novel treatment. The folate cycle containing several enzymes was identified as promising target; some of them are already addressed by antimalarials.^[1] However, inhibition of serine hydroxymethyl transferase (SHMT), a key enzyme of the folate cycle, has not been investigated so far. *A. Thaliana* SHMT inhibitors, based on a pyrazolopyran scaffold, from an herbicide optimization program at BASF-SE demonstrated promising antimalarial activity on *P. falciparum* and *P. vivax*.^[2] Because of pharmacokinetic limitations no significant activity in the *P. berghei* mouse model could be achieved.^[3] The binding mode was resolved by several X-ray crystal structures of PvSHMT-ligand complexes.^[3] Based on the high similarity of *P. vivax* and *P. falciparum* SHMT, the X-ray co-crystal structures can be utilized for 3D modeling to design small drug-like molecules against PfSHMT. In this work the development of novel inhibitors is focused on improving selectivity and liver microsomal stability while keeping high potency. In that perspective, variation of the moieties on the pyrazolopyran core and derivatization of the exit-vector are investigated.



[1] M. Schlitzer, *Arch. Pharm. Chem. Life Sci.* **2008**, 341, 149–163.

[2] M. Witschel, F. Stelzer, J. Hutzler, T. Qu, T. Mietzner, K. Kreuz, K. Groijmann, R. Aponte, H. W. Hoffken, F. Calo, T. Ehrhardt, A. Simon, and L. P. Rapado, *WO 2013/182472 A1* **2013**.

[3] M. Witschel, M. Rottmann, A. Schwab, U. Leartsakulpanich, P. Chitnumsub, M. Seet, S. Tonazzi, G. Schwartz, F. Stelzer, T. Mietzner, C. McNamara, F. Thater, C. Freymond, A. Jaruwat, C. Pinthong, P. Riengrunroj, M. Oufir, M. Hamburger, P. Mäser, L. M. Sanz-Alonso, S. Charman, S. Wittlin, Y. Yuthavong, P. Chaiyen, and F. Diederich, Inhibitors of Plasmodial Serine Hydroxymethyltransferase (SHMT): Cocrystal Structures of Pyrazolopyrans with Potent Blood- and Liver-Stage Activities, *J. Med. Chem.* **2015**, 58, 3117–3130.

Investigation of an engineered AaLS-13 capsid and Identification of the encapsulation pathway for GFP (+36) by high mass MALDI MS analysis

K. Root¹, R. Frey¹, D. Hilvert¹, R. Zenobi^{1*}

¹Department of Chemistry and Applied Biosciences, ETH Zurich, CH-8093 Zurich (Switzerland)

Proteins that self-assemble into nanocontainers are of emerging scientific interest for drug development.

Here, an engineered variant of Lumazine synthase from *Aquifex aeolicus* (AaLS-13) that self-assembles into a capsid was studied. Coulomb attraction between supercharged GFP (+36) and the capsid modified with additional negative charges on the lumen favors cargo encapsulation. We report on the successful implementation of native ESI-MS and MALDI-MS, to map the subunit composition at high molecular weight. The subunit stoichiometry of the AaLS-13 capsid was determined to be composed of only pentamer subunits. The capsid structure of AaLS-13 proposed earlier by Wörsdörfer *et al.* organizing into a regular icosaheder in the T=3 state, could not be confirmed, due to additionally required hexamer subunits [1]. Our results suggest the protein to build up a regular T=7d icosaheder capsid. High-mass MALDI-MS combined with crosslinking with glutaraldehyde (GA) is suggested to monitor the encapsulation pathway of supercharged GFP(+36) into the capsid. The results obtained showed capsid stabilization upon addition of supercharged GFP. Further, encapsulation of GFP (+36) strongly shifts the equilibrium towards capsid formation. The time scale of this reaction was determined to be out of range for crosslinking followed by high-mass MALDI analysis.

[1] B. Wörsdörfer, K. Woycechowsky, D. Hilvert, *Science*, **2011**, 331, 589-592.

Fluorine Scan at the Active Sites of Rhodesain and Human Cathepsin L: Enhanced Binding Affinity by Stacking of Fluorinated Phenyl Rings on Flat Dipeptide FragmentsM. Giroud¹, M. Harder¹, B. Kuhn², W. Haap², T. Schirmeister³, F. Diederich^{1*}¹ETH Zurich, ²F. Hoffmann-La Roche AG, ³University of Mainz

Amide bonds are ubiquitously present in proteins. However, exploitation of their planar π -surface for ligand binding remains to be further developed.^[1] Harder *et. al* showed that for efficient ligand–amide bond interactions (1) favorable dipole moment orientation between various heterocycles and *N*-methylacetamide (antiparallel alignment) and (2) reduced electron density of the heterocycle were required.^[2] Because of its high electronegativity and low polarizability, fluorine represents an ideal tool to (1) tune the electron density of benzene derivatives without introducing steric hindrance upon H/F replacement, and (2) introduce large dipole moment variations. Following these guidelines, we performed a fluorine scan of 20 inhibitors for two cysteine proteases, rhodesain and human cathepsin L, which both feature a flat dipeptide moiety in their respective pockets. We identified that multiple fluorination of the inhibitors led to an increased binding affinity for both rhodesain and human cathepsin L, resulting in low nanomolar inhibitors. Binding affinities were determined by fluorescence-based assays. We believe that these findings will not only be helpful to guide future development of inhibitors of rhodesain, a target against human African trypanosomiasis, but also to any other medicinal chemistry program whose target possesses exposed amide bonds.



[1] S. Cotesta, M. Stahl, *J. Mol. Model.* **2006**, 12, 436–444.

[2] M. Harder, B. Kuhn, F. Diederich, *ChemMedChem* **2013**, 8, 397–404. br

O⁶-Alkylguanine Post-lesion DNA synthesis by Y-family DNA polymerase ζ characterized with synthetic nucleosides

M. H. Rätz¹, H. Gahlon¹, L. Müller¹, S. Sturla^{1*}

¹Department of Health Sciences and Technology, Schmelzbergstrasse 9, 8092 Zürich, Switzerland

DNA polymerase ζ (Pol ζ) is a low-fidelity B-family polymerase involved in DNA damage tolerance and genomic stability by catalyzing the synthesis of DNA in the presence of DNA adducts. Pol ζ can bypass such chemical damage to the structure of DNA bases, but is in most cases inefficient at inserting a base opposite the adduct. In contrast, it seems to be very effective in performing post-lesion DNA synthesis (PLS), which is the extension step immediately following lesion bypass. Thus, Pol ζ may contribute to mutagenesis by fixing mismatches created by error-prone insertion.

In an effort to understand the mechanism of PLS for alkylated DNA of relevance to carcinogens that induce O⁶-alkylguanine adducts, which are known to induce mutations, we used adduct-directed synthetic nucleoside probes Benzi and BIM (Figure 1) with altered size and hydrogen bonding possibilities to test the structural requirements for Pol ζ PLS. Rates were characterized for DNA synthesis of terminal primer template mismatches comprised of G, O⁶-methylguanine (O⁶-MeG) and O⁶-(carboxymethyl)guanine (O⁶-CMG) paired opposite either canonical nucleotides or synthetic bases. When the adduct was paired at the template:primer terminus with a synthetic base with correct H-bond donor alignment with the adduct, extension was similar to canonical mismatch variations, whereas when the paired base had no minor-groove H-bond donor, extension was hindered. The rates of PLS were related with biophysical properties of the duplexes, and lesion/mismatch extension was found to be slowed in the presence of modification relative to an undamaged construct. These data provide new insight regarding contributions of hydrogen bonding interactions to promoting extension past O⁶-MeG and O⁶-CMG catalyzed by Pol ζ .





Figure 1. Hydrogen bonding capacity for nucleoside probes paired opposite O⁶-MeG (A) BIM opposite O⁶-MeG and (B) Benzi opposite O⁶-MeG.

Structural Characterization of OligoprolineP. Wilhelm¹, H. Wennemers^{1*}¹ETH Zurich, Laboratory of Organic Chemistry, Zurich, Switzerland

Oligoprolines have found wide applications as molecular scaffolds in various fields of chemistry and biology due to their well-defined and rigid conformation. In aqueous environments oligoprolines adopt a left-handed Polypyrrolone II (PPII)-helix with every third residue stacked on top of each other. This secondary structure is the third most common in proteins and is widely adopted by collagen, the most abundant protein in mammals.

Herein, we present a detailed analysis of the first high-resolution crystal structure of an oligoproline adopting a PPII-helix.





The crystal structure provided deep insights into the structural parameters of the PPII-helix. It revealed that neighboring amide bonds interact with each other and that water is not a prerequisite to form a PPII-helix.

Furthermore we determined the persistence length of oligoproline using EPR-spectroscopy. The obtained data are crucial for the use of oligoprolines as molecular scaffolds.

P. Wilhelm, B. Lewandowski, N. Trapp and H. Wennemers, *J. Am. Chem. Soc.*, **2014**, 136, 15829.
L. Garbuio, B. Lewandowski, P. Wilhelm, L. Ziegler, M. Yulikov, H. Wennemers, G. Jeschke, *Chem. Eur. J.*, **2015**, in press.

Cell Penetrating Peptides Based on an Oligoproline ScaffoldP. Raschle¹, H. Wennemers^{1*}¹ETH Zurich, Laboratory of Organic Chemistry, Zurich, Switzerland

The efficient uptake of drugs, biomolecules and imaging agents into cells is still limited by the poor translocation through the plasma membrane of eukaryotic cells. Cell Penetrating Peptides (CPPs) based on cationic and amphipilic oligoproline structures are promising tools to address this issue.^[1,2] To exploit the unique properties of oligoprolines as rigid and functionalizable scaffolds (Polyproline II helix)^[3] we designed well-defined guanidine decorated oligoprolines and explored their cell permeability properties. Oligoprolines with varying chain lengths, charge densities, as well as helicities were prepared. Their uptake into human cervical cancer (HeLa) cells was evaluated by flow cytometry. The quantitative uptakes of the peptides were compared to established CPPs like the Tat-peptide and oligoarginines. The intracellular localization of the peptides was analyzed by confocal microscopy. Additionally the stability of the oligoprolines was tested in human blood serum and trypsin.



[1] M. Zorko, Ü Langel, *Advanced Drug Delivery Rev.* **2005**, 57, 529.

[2] a) J. Fernandez-Carneado, M. J. Koagan, S. Castel, E. Giralt, *Angew. Chem. Int. Ed.* **2004**, 43, 1811. b) Y. A. Fillon, J. P. Anderson, J. Chmielewski, *J. Am. Chem. Soc.* **2005**, 127, 11798.


[3] a) K. Kuemin, S. Schweizer, C. Ochsenfeld, H. Wennemers, *J. Am. Chem. Soc.* **2009**, 131, 15474. b) M. Kümin, L.-S. Sonntag, H. Wennemers, *J. Am. Chem. Soc.* **2007**, 129, 566.

Oligoprolines as Scaffolds for Tumor Targeting with Hybrid Bombesin Analogues

S. Dobitz¹, C. Kroll¹, R. Mansi², B. Braun², H. R. Mäcke², H. Wennemers^{1*}

¹ETH Zurich, Laboratory of Organic Chemistry, Zurich, Switzerland, ²University Hospital Freiburg, Department of Nuclear Medicine, Freiburg, Germany

In aqueous environments oligoprolines adopt the well-defined polyproline II helix already at short chain lengths.^[1] Incorporation of 4-azidoproline residues into the sequence provides reactive sites in defined distances that can easily be functionalized through, e.g., click chemistry.^[2]



Hybrid ligands consisting of an oligoproline scaffold equipped with a bombesin-based agonist and antagonist exhibit extraordinary tumor uptake properties in prostate carcinoma.^[3] The hybrid ligands showed significantly higher uptakes *in vitro* and *in vivo* compared to monovalent and divalent controls. The defined distance between the recognition motives proved to be important for high, specific, and long lasting uptakes. Based on these initial findings we are now designing modified ligands to achieve higher uptakes and a deeper understanding of how the uptake is accomplished on the molecular and cellular level.

[1] a) F. Rabanal, L. M. Pons, E. Giralt, *Biopolymers* **1993**, 33,1019. b) S. Kakinoki, Y. Hirano, M. Oka, *Polym. Bull.* **2005**, 53, 109.

[2] a) M. Kümin, L.-S. Sonntag, H. Wennemers *J. Am. Chem. Soc.* **2007**, 129, 466. b) R. S. Erdmann, M. Kümin, H. Wennemers *Chimia* **2009**, 64, 19. c) Y. A. Nagel, M. Kümin, H. Wennemers *Chimia* **2011**, 65, 264.

Impact of minor groove alkylation on transcription by RNA polymerase II


S. Malvezzi¹, T. Angelov¹, L. Farnung², F. Eichenseher¹, P. Cramer², S. Sturla^{1*}

¹Health Sciences and Technology, ETH Zurich, ²Max Planck Institute for Biophysical Chemistry, Gottingen

Minor-groove binding chemicals can alkylate the 3-position of Adenosine, giving rise to adducts such as 3-methyl-adenine (3MeA), which is the major DNA adduct from methyl lexitropsin, an alkylating agent with potential utility in the therapy of gliomas. The effective cytotoxicity of minor groove alkylation may be reduced, however, by repair mechanisms like transcription-coupled nucleotide excision repair (TC-NER), which is activated when RNA polymerase II (RNAPII) stalls at DNA damage sites. In this study we describe the behavior of RNAPII in the presence of modified DNA including a depurination-resistant analog of 3MeA, 3-deaza-3-methyl-adenine (3dMeA). We performed transcription assays with human RNAPII derived from HeLa nuclear extract and primer extension of ternary elongation complexes with purified yeast RNAPII. Our results show that although RNAPII efficiently synthesizes past 3dMeA, a significant reduction in the formation of full-length product was observed in comparison to transcription of an unmodified DNA template. Moreover, RNAPII promoted the mutagenic incorporation of cytosine opposite 3dMeA, a phenomenon that may lead to transcriptional mutagenesis in cells. Additional data from studies concerning how steric and H-bonding properties of minor groove modifications modulate these processes will be presented. Taken together, these results provide new chemical insight concerning how RNAPII copes with DNA alkylation, and this information is relevant for how tumor cells may resist chemotherapy.

Novel azobenzene-derived visible light photoswitches for biological applicationsZ. Pianowski¹, M. Steinbiss^{1,2}, J. Karcher¹¹Karlsruher Institut für Technologie (KIT), ²Currently: TU Braunschweig

Molecular photoswitches are compounds that undergo reversible structural changes upon irradiation with light. After covalent or non-covalent attachment to biopolymers they enable photomodulation of their structure or activity. Azobenzene derivatives are most common molecular photoswitches applied in biological systems. However, unmodified azobenzene is switched with UV light, insoluble in water and moderately resistant on intracellular reducing environment. Therefore structural modifications are developed to increase the scope of applications, especially inside living cells and organisms.




We have synthesized several arylazopyridinium salts which absorb in the visible wavelength range, are well-soluble in aqueous buffers and tolerate glutathione at physiological concentrations. We will present our most recent results and discuss the scope of applications of our compounds for biological systems.

Bacterial Resistance to Silver: The Role of SilE ProteinV. Chabert¹, K. Fromm^{1*}¹University of Fribourg

Silver has been used for hundreds of years for its antimicrobial properties. Since the emergence of many multi-resistant bacterial strains against classical antibiotics, the research of new silver compounds is now at its apogee. Nowadays, a lot of researches are focused on compounds with slow- and stimuli-responsive- release of Ag^+ . While these drugs have been shown to be highly able to kill bacteria, some of these pathogens have developed a resistance to high concentrations of Ag^+ .

This resistance is provided by the plasmid pMG101, which encodes for eight proteins that act together in an efflux pump system to deal with silver ions. Among these, the SilE protein is the only one of which its mode of action is actually unknown.



To identify the role of SilE in this bacterial machinery, two approaches have been intended in our group. While one way is to study the interaction of the whole protein with silver ions, the other is based on a bottom-up approach, investigating the interaction of silver ions with short peptide sequences of this protein.

S. Silver, L. T. Phung, G. Silver, *J. Ind. Microbiol. Biotechnol.* **2006**, 33, 627-634.

S. Eckhardt, P. S. Brunetto, J. Gagnon, M. Priebe, B. Giese, K. M. Fromm, *Chem. Rev.* **2013**, 113, 4708-4754.

The use of phosphorylated peptides to explore the folding properties of the protein tau required for AT8 antibody recognition

Y. Jacquot¹, F. Cantrelle², I. Landrieu², C. Byrne¹, K. Pedrag³, L. Amniai², J. Wieruszeski², R. L. Mancera⁴, G. Lippens^{2*}

¹Université Pierre et Marie Curie, Paris, ²University of Lille1, France, ³University of Cambridge, UK, ⁴Curtin University, Perth, Australia

Paired helical filaments (PHFs) resulting from the hyperphosphorylation of the protein tau are major components of the Alzheimer disease. They contribute to the formation of intraneuronal neurofibrillary tangles. Accordingly, the phosphorylation of the serine 202 (Ser-202) and the threonine 208 (Thr-208) seem relatively specific. These two residues are, indeed, key for the recognition of the AT8 antibody [1], which is widely used for post mortem diagnosis of Alzheimer's disease. Although it is well established that the epitope of AT8 is located around the Ser-202 and the Thr-205, the folding characteristics associated with the recognition of this region with the AT8 are not known.

By using NMR spectroscopy in combination with scaled molecular dynamics (SMD) on different phosphorylated peptides corresponding to the 192-212 fragment of tau, we studied the secondary structure of the AT8 epitope of tau under its hyperphosphorylated form [2]. We observed that the AT8 epitope forms a turn that is induced by the phosphorylated threonine 205 (pThr-205) and the glycine 207 (Gly-207). More precisely, we detected hydrogen interactions between the phosphate group of the pThr-205 and the amide proton of the Gly-207. This interaction was stabilized by electrostatic contacts between the pSer-202/pThr-205 side chains and the Arg-209/Arg211. These results were compared with those obtained from the hyperphosphorylated full-length version of tau obtained by using the recombinant CDK2/CycA3 kinase [2].

[1] M. Goedert, R. Jakes, E. Vanmechelen. Monoclonal antibody AT8 recognises tau protein phosphorylated at both serine 202 and threonine 205. *Neurosci. Lett.* **1995**, 189, 167-169.

[2] N.S. Gandhi, I. Landrieu, C. Byrne, P. Kukic, L. Amniai, F.X. Cantrelle, J.M. Wieruszeski, R.L. Mancera, Y. Jacquot, G. Lippens. A phosphorylation-induced turn defines the Alzheimer's disease AT8 antibody epitope on the tau protein. *Angew. Chem. Int. (Ed. English)* **2015** (in press).b

Pollen induced asthma - could small molecules in pollen exacerbate the protein-mediated allergic response?A. Bozicevic¹, M. De Mieri¹, M. Hamburger^{1*}¹University of Basel

Plant pollen are known to be strong airborne elicitors of asthma in humans. *In vitro* data and clinical studies corroborate the involvement of small surface proteins present on the pollen grain. They modulate the immune system through IgE cross-linkage causing airway inflammation, and obstruction due to constriction of airways. At the physiological level, relaxation and constriction of airways is regulated by mechanisms involving proteins such as the lipid kinase PIP5K γ and the cation channel TRPA1. While the role of proteins is well established, a possible contribution of small molecules present in pollen to the clinical outcome of asthma has not been explored up to now. Therefore, we analyzed and compared the phytochemical profiles of pollen originating from thirty plant species causing varying degrees of pollen allergenicity. Profiling was performed with high performance liquid chromatography (HPLC) coupled with electrospray ionization mass spectrometry (ESIMS), photodiode array (PDA) and evaporative light scattering (ELSD) detectors, and supported by microprobe nuclear magnetic resonance (NMR) spectroscopy and spectrophotometric analysis. The presence of conjugated polyamines, such as N¹,N⁵,N¹⁰-tricoumaroylspermidine, N¹-caffeoyl-N⁵,N¹⁰-dicoumaroylspermidine and N¹,N⁵,N¹⁰,N¹⁵-tetracoumaroylspermine was a characteristic feature of pollen from Asteraceae (*Ambrosia* and *Artemisia* ssp.), and compounds with Michael acceptor properties were also mainly present in pollen of these species. Polyamines such as spermine and spermidine are modulators of the lipid kinase PIP5K γ ^[1], and sesquiterpene lactones activate cation channel TRPA1^[2]. Thus, the possible contribution of these small molecules in the exacerbation of airway constriction should be explored in more detail.

[1] Erle and Sheppard, *J. Cell Biol.* **2014**, 205, 621–631.

[2] Avonto et al., *Angew. Chem. Int. Ed.* **2011**, 50, 467–471.

Search for alternatives to copper in organic farming: Fungicidal activity of a *Juncus effusus* medulla extract and its active constituent, dehydroeffusol, against downy mildew and apple scab


J. Ramseyer¹, B. Thuerig², M. De Mieri¹, F. Schmidt¹, O. Potterat¹, H. Schärer², T. Oberhänsli², L. Tamm², M. Hamburger^{1*}

¹University of Basel, ²Research Institute of Organic Agriculture (FiBL)

Copper has been used since the 19th century for the control of plant diseases, and is still permitted in organic agriculture out of this tradition. In recent years, the utilization of copper has been criticized due to an unfavourable ecotoxicological profile [1]. Even though the amounts of copper have been significantly reduced, copper input is still above uptake by plants, and results in accumulation in the soil. Therefore, considerable efforts have been made in organic agriculture to identify ecologically safer substitutes.

In this context, an in-house library of plant and fungal extracts was screened *in vitro* for an inhibitory effect against several plant pathogens (fungi, oomycetes, bacteria) [2]. Hits were further assessed on grapevine and apple seedlings. Active constituent(s) were identified by a procedure referred to as HPLC-based activity profiling which combines biological activity data with chemoanalytical information. Structure elucidation was performed by a combination of ESI-MS and NMR spectroscopy.

As one of the hits, the ethyl acetate extract of *Juncus effusus* L. (Juncaceae) medulla showed strong inhibitory activity against *Venturia inaequalis* (apple scab) and *Plasmopara viticola* (grapevine downy mildew), with mean minimal inhibitory concentrations (MIC) (100%) of 35 µg/mL and 25 µg/mL, respectively. In a secondary assay on grapevine leaf discs inoculated with *P. viticola*, 94% inhibition was observed at a concentration of 0.5 mg/mL. When tested on grapevine and apple seedlings at a concentration of 0.5 mg/mL, the growth of these fungi was, on average, inhibited with 98% and 84% efficacy, respectively. The active constituent was identified as dehydroeffusol (**1**) [3], and showed mean MICs of 12 µg/mL against *V. inaequalis*, and 4.1 µg/mL against *P. viticola*, *in vitro*. Subsequent *in vivo* assessment of the pure compound revealed inhibition rates of 82% on grapevine seedlings, and 86% on apple seedlings at a concentration of 32 µg/mL. Our results demonstrate that plants can provide promising opportunities for the replacement of copper in organic farming.



[1] Christine Kula, Susanne Guske, *Berichte aus der biologischen Bundesanstalt für Land- und Forstwirtschaft*, **2003**, 11-16.

[2] Olivier Potterat, Matthias Hamburger, *Planta Medica*, **2014**, 80, 1171-1181.


[3] Katsuhito Shima, Masao Toyota, Yoshinori Asakawa, *Phytochemistry*, **1991**, 30, 3149-3151.

Natural and semisynthetic antitrypanosomal sesquiterpene lactones from *Anthemis nobilis*

M. De Mieri¹, I. Ismajili¹, M. Kaiser², R. Brun², M. Hamburger^{1*}

¹University of Basel, ²Swiss Tropical and Public Health Institute, Basel

Trypanosomatid parasites of the genera *Trypanosoma* and *Leishmania* are responsible for neglected tropical diseases for which new, safer, and affordable drugs are urgently needed. [1] Natural products from plants have been instrumental in developing drugs against protozoal diseases, e.g. artemisinin and quinine, and still represent an important source for leads with trypanocidal activity. [2] In previous studies sesquiterpene lactones (STLs) were found to possess promising antitrypanosomal activity. [3] We recently reported on the synthesis, by biomimetic acid catalysis, of a focused library of antitrypanosomal compounds generated from nobilin (**1**), the major STL in Roman chamomile (*Anthemis nobilis*) flower cones. [4] We here applied the same biomimetic approach to the 1(10) and 4(5) epoxides of **1** (**2** and **3**) and obtained a series of furanoheliangolide STL derivatives (**4-7**). Moreover, further phytochemical investigation of *A. nobilis* afforded a series of minor STLs, four of which (**8-11**) were new compounds. All the above mentioned STLs showed high antitrypanosomal activity (IC_{50} s 0.13-0.82 $\pm$ 0.01 μ M and 0.14-10.85 $\pm$ 0.01 μ M against *T. b. rhodesiense* and *L. donovani*, respectively), although with low selectivity. SAR studies are currently ongoing to identify structural and physico-chemical features responsible for activity.



[1] K.T. Andrews et al. *Int. J. Parasitol. Drugs Drug Resist.* **2014**, 4, 95-111.

[2] T. Schmidt et al. *Curr. Med. Chem.* **2012**, 19, 2128-2175.

[3] S. Zimmermann et al. *Planta Med.* **2012**, 78, 553-556.

[4] M. De Mieri et al. *Bioorg. Med. Chem.* **2015**, 23, 1521-1529.

Antagonizing Bacterial Adhesion - Hit Identification by a Dynamic Combinatorial Chemistry Approach

P. Frei¹, L. Pang¹, D. Eris¹, B. Ernst^{1*}

¹University of Basel

In Dynamic Combinatorial Chemistry (DCC) substance libraries are formed from reversibly reacting building blocks. These libraries are under thermodynamic control, and their composition can be influenced by the applied conditions. For instance, in the presence of a receptor protein, those members of the library binding to the target are up-regulated, leading to a shift of the composition of the library, i.e. an amplification of high-affinity ligands at the expense of weak binders.^[1,2]

Urinary tract infections belong to the most common bacterial infections and are mainly caused by uropathogenic *Escherichia coli* (UPEC). The infection cycle is initiated by the binding of UPEC via the lectin FimH to uroplakin 1a, a mannosylated glycoprotein on bladder epithelial cell.^[3,4] This interaction can be prevented with aryl mannosides leading to elimination of bacteria with the bulk flow of urine and thus presenting an alternative treatment strategy to classical antibiotics.^[5-8]

We here present a DCC approach for the bacterial lectin FimH. Aryl glycosides with an aldehyde moiety in the aglycone are reacted reversibly with hydrazides to generate a library of acylhydrazones. The synthesis is carried out in the absence and presence of the FimH. The library is then rendered pseudostatic by increasing the pH, making it suitable for HPLC analysis. Comparing the library compositions obtained in the presence and in absence of the lectin allows for quantitative affinity ranking of the corresponding acylhydrazones.

[1] O. Ramström, J.-M. Lehn, *Nat. Rev. Drug. Discov.* **2002**, 1, 26.

[2] M. Mondal, A.K.H. Hirsch, *Chem. Soc. Rev.*, **2015**, 44, 2455.

[3] T.J. Wiles, R.R. Kuluesus, M.A. Mulvey, *Exp. Mol. Pathol.* **2008**, 85, 11.

[4] J.D. Schilling, M.A. Mulvey, S.J. Hultgren, *J. Infect. Dis.* **2001**, 183 Suppl. 1, S36.

[5] O. Schwardt *et al.* *Bioorg. Med. Chem.* **2011**, 12, 6454.

[6] L. Pang *et al.* *ChemMedChem* **2012**, 7, 1404.

[7] X. Jiang *et al.* *J. Med. Chem.* **2012**, 55, 4700.

[8] S. Kleeb *et al.* *J. Med. Chem.* **2015**, in press.

It's better to bend than to break

S. Rabbani¹, P. Zihlmann¹, B. Fiege¹, C. Sager¹, R. A. Jacob², R. Vannam³, M. Peczu³, B. Ernst^{1*}

¹University of Basel, Molecular Pharmacy, ²University of Basel, Structural Biology, ³University of Connecticut, Department of Chemistry

The attachment of uropathogenic *E. coli* to host cells is mediated by the adhesin FimH which is located at the tip of bacterial type 1 pili. The carbohydrate recognition domain of FimH forms an extended hydrogen bond network with the hydroxyl groups of aryl α -D-mannosides. In addition, the so-called tyrosine-gate hosts the aryl aglycone.

Herein we describe the molecular interaction between flexible seven membered ring (septanose) monosaccharides that deform upon binding to the rigid FimH to accommodate to the shape of the binding pocket. The interaction was firstly evaluated in a competitive binding assay and molecular docking experiments. Subsequent isothermal titration calorimetry (ITC) measurements, molecular dynamics (MD) simulations and NMR chemical shift perturbation (CSP) experiments revealed that the difference in affinity between *n*-heptyl α -D-mannoside and the corresponding septanose derivative was solely due to an induced conformational change (entropy cost) of the septanose ligand. Finally, the cocrystal structure confirmed the binding mode of the septanose to FimH.

Combining different analytical methods considerably improved our understanding of FimH-ligand interaction and will guide the design of therapeutically relevant FimH antagonists.

Keywords: FimH, mannosides, NMR, molecular dynamics, thermodynamics, X-Ray.

Development of small molecular tools for the cellular study of adenosine A₁ receptors


J. Hemmings¹, A. Knight², B. Frenguelli², S. J. Dowell³, G. Ladds⁴, M. Lochner^{1*}

¹University of Bern, ²University of Warwick, ³GlaxoSmithKline, ⁴University of Cambridge

The adenosine receptors are members of the G-protein coupled receptor (GPCR) family which represents the largest class of cell-surface proteins mediating cellular communication. As a result, GPCRs are formidable drug targets and it is estimated that approximately 30% of the marketed drugs act through members of this receptor class. There are four known subtypes of adenosine receptors: A₁, A_{2A}, A_{2B} and A₃. The adenosine A₁ receptor, which is the subject of this presentation, mediates the physiological effects of adenosine in various tissues including the brain, heart, kidney and adipocytes. In the brain for instance, its role in epilepsy and ischemia has been the focus of many studies.^[1]

Previous attempts to study the biosynthesis, trafficking and agonist-induced internalisation of the adenosine A₁ receptor in neurons using fluorescent protein-receptor fusion constructs^[2] have been hampered by the sheer size of the fluorescent protein (GFP) that ultimately affected the function of the receptor. We have therefore initiated a research programme to develop small molecule fluorescent agonists that selectively activate the adenosine A₁ receptor.

Our probe design is based on the endogenous ligand adenosine and the known unselective adenosine receptor agonist NECA (Figure). We have synthesised a small library of non-fluorescent adenosine derivatives that have different cyclic and bicyclic moieties at the N⁶ position of the purine ring and have evaluated the pharmacology of these compounds using a yeast-based assay.^[3] This analysis revealed compounds with interesting behaviour, i.e. exhibiting adenosine A₁ subtype-selectivity or G-protein bias. Thus, these compounds can be potentially used as tool compounds in their own right for cellular studies of the adenosine A₁ receptor. Furthermore, we have also linked fluorescent dyes to the purine ring and discovered fluorescent compounds that can activate the adenosine A₁ receptor.



[1] N. Dale, B. N. Frenguelli, *Curr. Neuropharm.* **2009**, 7, 160-179.

[2] A. E. Baines, S. A. L. Correa, A. J. Irving, B. G. Frenguelli, *Neuropharmacology* **2011**, 61, 1-11.

[3] G. D. Stewart, C. Valant, S. J. Dowell, D. Mijaljica, R. J. Devenish, P. J. Scammells, P. M. Sexton, A. Christopoulos, *J. Pharmacol. Exp. Ther.* **2009**, 331, 277-286.

Fluorescent probes for the cellular study of the 5-HT₃A receptor - synthesis and evaluation of near-infrared probes

J. Simonin¹, T. Jack¹, M.-D. Ruepp¹, J. Gertsch¹, A. J. Thompson², M. Lochner^{1*}

¹University of Bern, ²University of Cambridge

The 5-HT₃ receptor is a ligand-gated ion channel (LGIC) and a member of the Cys-loop family of receptors. Fluorescent probes are powerful tools to study transmembrane proteins such as the 5-HT₃. We previously reported^{[1],[2]} a fluorescein derivative that showed high affinity (1.9 nM) and that was used as a tracer in fluorescence polarization and flow cytometry binding assays.

A near-infrared fluorescent probe was used for in-vivo imaging of the 5-HT₃R in a live mouse^[2] and stained receptors in the abdominal cavity, in intestines, but also in salivary glands. The unexpected presence of 5-HT₃ receptors in mouse salivary glands was confirmed by Western blots. This probe was also used in single molecule spectroscopy studies of the receptor^[3]. Herein we report a straightforward synthesis of the SiRhodamine methyl and lactone fluorophores their live cells evaluation and the evaluation of their granisetron conjugates.




Fig. 1. 10 mg/kg i.p. of G-SiR (640/680 nm) life imaging with anesthetized mice, 5 minutes post injection and 30 minutes post injection.

[1] J.Simonin; S.K.V.Vernekar; A.J.Thompson; J.D. Hothersall; C.N. Connolly; S.C.R.Lummis; M.Lochner, *Bioorg. Med. Chem. Lett.*, **2012**, 22, 1151.

[2] T.Jack; J.Simonin; M.-D.Ruepp; A. J.Thompson; J.Gertsch; M.Lochner, *Neuropharmacology*, **2015**, 90, 63-73.

[3] Adam O. Barden et al., *Neuropharmacology*, manuscript accepted.

Peptide dendrimer as SiRNA transfection reagentM. Heitz¹, T. Darbre^{1*}, J.-L. Reymond^{1*}¹University of Bern

RNA interference (RNAi) allows effective and specific silencing as described by Tuschl and co-workers in their proof-of-principle experiment demonstrating that synthetic double stranded small interfering RNA (SiRNA) could achieve sequence-specific gene knockdown in a mammalian cell line by promoting the degradation of complementary mRNA via RISC complex¹. Potential therapeutic applications of RNAi are crucially dependent on the delivery of SiRNA into the cytosol to avoid that this step becomes a bottleneck. Naked or chemically modified SiRNA delivery is of limited application and therefore nanoparticles encapsulating SiRNA molecules have been investigated as a more general method to bring SiRNA into cells.

We have previously explored a collection of peptide dendrimers for the transfection of plasmid DNA and found efficient reagents that obeyed structure-activity relationships. Of crucial importance was the distribution of cationic charges across the three dendrimer generations and the use of DOTMA/DOPE as lipids (Figure 1, A)².




Figure 1. The chemical structure of the peptide dendrimer G3-KL1,2,3 used in DNA transfection (A). SiGAPDH expression following a transfection in presence of 10 % serum of the lead compounds and lipofectamine 2000 in HeLa cells compared to untreated cells (B).

We are now exploring peptide dendrimers as delivery agents for SiRNA. In this project a library of 100 peptide dendrimers were prepared by SPPS and their gene silencing ability investigated. The parameters necessary for efficient gene silencing have been discovered and optimized to lead to an only amino acid and a lipid-conjugated dendrimer (Figure 1, B). The biological experiments included treatment of HeLa cells by the new transfection agents and SiRNA targeting GAPDH (SiGAPDH) or scrambled (SiNC) in the absence and presence of serum. The knockdown efficiency was measured by monitoring enzyme activity of GAPDH.

[1] Sayda M. Elbashir, Jens Harborth, Winfried Lendeckel, Abdullah Yalcin, Klaus Weber, Thomas Tuschl, *Nature*, 2001, 411, 494-498.


[2] Albert Kwok, Gabriella Eggimann, Jean-Louis Reymond, Tamis Darbre, Florian Hollfelder, *ACS Nano*, 2013, 7, 4668-4682.

Excess Electron Transfer in DNA Containing a Pyrenyl Donor and Multiple Stable Phenanthrenyl Base-Surrogates

P. R  thlisberger¹, C. Leumann^{1*}

¹Universit  t Bern

The reductive electron transfer along DNA attracted considerable interest in the last two decades and is a promising feature for molecular electronics. A system based of a p-stacked C-nucleoside pyrenyl donor (C-AP), multiple phenanthrenyl based base surrogates as electron carriers and 5-bromodeoxyuracil (BrdU) as kinetic electron trap was synthesized in order to study the excess electron transfer (EET). Gradually lowering the lowest unoccupied molecular orbital (LUMO) of the sequentially installed electron carriers is envisioned to enhance the electron transfer efficiency by disfavoring the back electron transfer.



- [1] N.A. Grigorenko, C.J. Leumann, *Chem. Eur. J.* **2009**, 15, 639.
[2] N.A. Grigorenko, C.J. Leumann, *Chem. Commun.* **2008**, 14, 5417.
[3] P. Roethlisberger, F. Wojciechowski, C.J. Leumann, *Chem. Eur. J.*, **2013**, 19, 11518-11521.

Antimicrobial Cyclic Peptides with L,D- architecture Targeting Pseudomonas aeruginosaR. He¹, T. Darbre¹, J.-L. Reymond^{1*}¹University of Bern


Membrane active antimicrobial peptides (AMP) with unusual topologies offer an attractive opportunity to address multi-antibiotic resistant bacteria, in particular Gram negative *Pseudomonas aeruginosa* (PA), which is one of the most problematic pathogens in the clinic.[1] Six and eight residue cyclic D,L- α -peptides have been reported to show antimicrobial activities against Gram positive strains such as *S. aureus* [2]. We asked the question whether this architecture could also be used to obtain AMPs targeting PA. A modified synthesis using a simple thioether ligation for cyclization was employed to prepare cyclic 6- to 12-residues peptides with alternating D- and L-residues, leading to the identification of two potent tryptophan containing cyclic peptides with strong activity against PA. CD spectroscopy shows that these active cyclic peptides undergo specific conformational changes in membrane environment compared to inactive analogs. Moreover, a red shift in the UV-VIS spectrum in SDS solution indicates that tryptophan residues form J-aggregates, suggesting aggregation in the membrane of bacteria as the mechanism of antibacterial action.

Reference:

- 1.Stach M, Reymond JL,et.al.*Angew Chem Int Ed Engl.***2014**,53, 12827-12831.
- 2.S. Fernandez-Lopez, M. Ghadiri,et al.*Nature.*,**2001**, 412, 452-455

Synthesis of a Geminal Difluorinated Tricyclic Nucleoside AnalogS. Frei¹, C. Leumann^{1*}¹University of Bern

In recent years fluorine substituted compounds have become particularly attractive in medicinal chemistry due to the bio-stability enhancing properties of the fluorine atom. [1] This properties do not only pertain to small molecules but also to oligonucleotides. This led us to synthesize and investigate therapeutic antisense oligonucleotides bearing a fluorine substituent at the C6' position of a bicyclic [3] or tricyclic [4] as well as at the C2' position [5] of a tricyclic nucleoside analog. In all cases these oligonucleotides were found to be good RNA mimics. Here we will show and discuss the synthesis of a C8' gem-difluoro modified tricyclic nucleoside analog. The key step in the synthetic pathway was the treatment of a bicyclic silyl enol ether with a difluoro carbene precursor reagent to form the gem-difluoro cyclopropane ring of the tricyclic sugar backbone with the desired exo-configuration.



[1] William K. Hagmann, *J. Med. Chem.*, **2008**, 51, 4359-4369

[2] Branislav Dugovic and Christian J. Leumann, *J. Org. Chem.*, **2014**, 79, 1271-1279

[3] Michal Medvecky, Alena Istrate, and Christian J. Leumann, *J. Org. Chem.*, **2015**, 80, 3556-3565

[4] Alena Istrate, Michal Medvecky, and Christian J. Leumann, *Org. Lett.*, **2015**, 17, 1950-53

Synthesis of photo-crosslinking probes and their application for the site-selective chemical modification of the 5-HT₃ receptor


T. Jack¹, M. Leuenberger¹, M.-D. Ruepp¹, S. Braga Lagache¹, O. Mühlemann¹, A. J. Thompson², M. Heller¹, M. Lochner^{1*}

¹University of Bern, ²University of Cambridge

The 5-HT₃ receptor (5-HT₃R) is an important ion channel responsible for the transmission of nerve impulses in the CNS and PNS and is activated by the endogenous agonist serotonin (5-hydroxytryptamine, 5-HT). The 5-HT₃R is the only serotonin receptor belonging to the Cys-loop superfamily of neurotransmitter receptors. Different structural biology approaches can be applied, such as crystallization and x-ray analysis.¹ However, characterizing the exact ligand binding site(s) of these dynamic receptors remains challenging. The use of photo-crosslinking probes is an alternative validated approach and allows identification of regions in the protein that are important for the binding of small molecules.

We designed our probes based on the core structure of the 5-HT₃R antagonist granisetron, an FDA approved drug used for the treatment of chemotherapy-induced nausea and vomiting.

We synthesized a small library of photo-crosslinking probes by conjugating diazirines and benzophenones via various linkers to granisetron. We were able to obtain several compounds with diverse linker lengths and different photo-crosslinking moieties that show nanomolar binding affinity for the orthosteric binding site. Furthermore we established a stable h5-HT₃R expressing cell line and a purification protocol to yield the receptor in high purity. Several experiments showed unambiguously that we are able to photo-crosslink our probes with the receptor site-specifically. The functionalised protein was analysed by Western blot and MS-analysis.² This yielded the exact covalent modification site, corroborating current ligand binding models derived from mutagenesis and docking studies.



[1] G. Hassaine *et al.*, *Nature* **2014**, 512, 276-281.

[2] T. Jack *et al.*, *in preparation*.

Design of 3D Protein Fingerprint and its Application to Map the Protein Data BankX. Jin¹, M. Awale¹, J.-L. Reymond^{1*}¹University of Bern

The RCSB Protein Data Bank (PDB) provides public access to experimentally determined 3D-structures of biological macromolecules (proteins, peptides and nucleic acids).¹ Although various tools are available to explore the PDB for analogs of specific proteins, the overall structural diversity available in PDB is difficult to perceive.²⁻⁶ Inspired by fingerprint based similarity searches which can be performed within seconds and greatly facilitate the exploration of small molecule databases,⁷ we have adapted our recently reported fingerprint-browser⁷ and fingerprint-mapplet⁸ concepts to enable the visualization and similarity search of PDB in form of a “PDB-Mapplet” application. PDB structures are encoded in a 136-bit atom pair 3D protein fingerprint (3DP) counting categorized atom pairs at increasing through-space distances.⁹ This fingerprint perceives molecular shape and groups PDB structures according to 3D-features such as folds rather than by primary sequence. The related 136-dimensional molecular shape space is visualized by principal component analysis of a secondary 200-dimensional satellite similarity space¹⁰ and representation of the (PC1,PC2)-plane. This plane covers 98 % of data variability and distributes PDB structures according to size and shape.

The PDB-Mapplet consists of a main window displaying interactive “google-maps-like” color-coded PC-maps, and locate or search for nearest neighbours is available for a given pdb entry. Each PDB structure is displayed as a cartoon image and can be inspected closer as 3D-model by opening a secondary Jmol visualisation window. The PDB-Mapplet offers an unprecedented insight into the diversity of PDB-structures and allows perceive relationships between proteins otherwise difficult to identify.

[1] Peter W. Rose, Andreas Prlić, Chunxiao Bi, Wolfgang F. Bluhm, Cole H. Christie, Shuchismita Dutta, Rachel Kramer Green, David S. Goodsell, John D. Westbrook, Jesse Woo, Jasmine Young, Christine Zardecki, Helen M. Berman, Philip E. Bourne, and Stephen K. Burley, *Nucleic Acids Res.*, **2015**, 43, (Database issue), D345-D356.

[2] Antonina Andreeva, Dave Howorth, Cyrus Chothia, Eugene Kulesha, and Alexey G. Murzin, *Nucleic Acids Res.*, **2014**, 42, (Database issue), D310-D314.

[3] CA Orengo, AD Michie, S Jones, DT Jones, MB Swindells, and JM Thornton, *Structure*, **1997**, 5, 1093-1108.

[4] Liisa Holm, and Päivi Rosenström, *Nucleic Acids Res.*, **2010**, 38, (Web Server issue), W545-W549.

[5] Ilya N. Shindyalov, Philip E. Bourne, *Protein Eng.*, **1998**, 11, 739-747.

[6] David La, Juan Esquivel-Rodríguez, Vishwesh Venkatraman, Bin Li, Lee Sael, Stephen Ueng, Steven Ahrendt, and Daisuke Kihara, *Bioinformatics*, **2009**, 25, 2843-2844.

[7] Mahendra Awale, and Jean-Louis Reymond, *Nucleic Acids Res.*, **2014**, 42, 234-239.


[8] Mahendra Awale, Ruud van Deursen, and Jean-Louis Reymond, *J. Chem. Inf. Model.*, **2013**, 53, 509-518.

[9] Mahendra Awale, Xian Jin, and Jean-Louis Reymond, *J. Cheminform.*, **2015**, 7, 3.

[10] José L. Medina-Franco, Gerald M. Maggiora, Marc A. Giulianotti, Clemencia Pinilla, and Richard A. Houghten, *Chem. Biol. Drug Des.*, **2007**, 70, 393-412.

QPD-BA, A Precipitating Dye For Monitoring Hydrogen Peroxide In Living CellsE. Lindberg¹, L. Holtzer¹, M. Gonzalez-Gaitan¹, N. Winssinger^{1*}¹University of Geneva

Reactive oxygen species such as hydrogen peroxide serve a variety of functions in living systems, including protein modification, signaling, immune and growth responses[1][2]. Several fluorescent probes selectively detecting hydrogen peroxide based on boronate oxidation have been reported [3].



Thus a quinazolinone precipitating dye (QPD) was modified with an immolative boronic acid trigger. Cleavage of the boronic acid moiety in the presence of hydrogen peroxide induced the precipitation of QPD which could be visualized in living cells.

[1] Michael Schieber, Navdeep S. Chandel, *Curr. Biol.*, **2014**, 24, R453-R462.

[2] Kira M. Holmström, Toren Finkel, *Nat. Rev. Mol. Cell Biol.*, **2014**, 15, 411-421.


[3] Alexander R. Lippert, Genevieve C. Van De Bittner, Christopher Chang, *Acc. Chem. Res.*, **2011**, 44, 793-804.

Protein Glycoconjugation by bioorthogonal click chemistry

T. Machida¹, K. Lang², L. Xue³, J. Chin⁴, N. Winssinger^{1*}

¹University of Geneva, ²Technical University Munich, ³Ecole Polytechnique Fédérale de Lausanne (EPFL), ⁴Medical Research Council

Protein glycosylation constitute an important post-translational modification and the glycans of glycoprotein have been specifically associated with numerous biological processes. We developed biocompatible methods to chemoselectively glycosylate protein in the lysate or live cell situation. Tetrazine-glycan conjugate allowed site-selectively glycosylation with genetically encoded *trans*-cyclooctene (TCO) protein in E coli pellet¹. Similarly, CuAAC based glycosylation proceeded in live cell under low copper concentration (50nM, 5 min reaction) in alkyne labeled Hela cell (metabolic labeling of homopropargylglycine)².



[1] K. Lang, L. Davis, S. Wallace, M. Mahesh, D. J. Cox, M. L. Blackman, J. M. Fox, J. W. Chin, *J. Am. Chem. Soc.* **2012**, 134, 10317–10320.

[2] C. Uttamapinant, A. Tangpeerachaikul, S. Grecian, S. Clarke, U. Singh, P. Slade, K. R. Gee, A. Y. Ting, *Angew. Chem. Int. Ed.* **2012**, 51, 5852–5856.


Platinum-Porphyrin Conjugates as Highly Phototoxic Agents against Human Cancer Cells

B. Spingler¹, A. Naik¹, R. Rubbiani¹, G. Gasser¹

¹University of Zurich

Despite the extended use of porphyrin complexes in PDT [1], tetraplatinated porphyrins have so far not been studied for their anticancer properties in the presence of light [2]. This is in contrast to ruthenated tetrapyrrolyl porphyrin complexes which have been synthesized [3] and tested for their PDT ability in animal models [4].

We would like to report about the synthesis of novel tetraplatinated porphyrins **1** as well as their photophysical characterization and *in vitro* light-induced anticancer properties [5]. The quantum yield of $^1\text{O}_2$ (Φ) production upon light irradiation was found to be between 0.42 and 0.54. The dark and light toxicity against human cancerous and non-cancerous cell lines (MRC-5, HeLa, A2780 and CP70) was determined by the resazurin assay. IC_{50} values were obtained after 4 h incubation, followed by 15 min irradiation at either 420 nm or 575 nm respectively. These platinum-porphyrin conjugates **1** had only minor dark toxicity, however upon visible light irradiation, IC_{50} values down to 19 ± 4 nM could be observed. These values correspond to an excellent phototoxic index ($\text{PI} = \text{IC}_{50} \text{ dark} / \text{IC}_{50} \text{ light}$) of greater than 5000. The influence of the absence / presence of different metals inside the porphyrin upon photophysical and pharmacological properties will be discussed as well.



1) MacDonald, I. J.; Dougherty, T. J. *J. Porphyrins Phthalocyanines* **2001**, 5, 105-129; Agostinis, P.; *et al. CA Cancer J. Clin.* **2011**, 61, 250-281.

2) Munakata, H.; Imai, H.; Nakagawa, S.; Osada, A.; Uemori, Y. *Chem. Pharm. Bull.* **2003**, 51, 614-615.

3) Gianferrara, T.; Bratsos, I.; Iengo, E.; Milani, B.; Ostric, A.; Spagnul, C.; Zangrando, E.; Alessio, E. *Dalton Trans.* **2009**, 10742-10756; Schmitt, F.; Govindaswamy, P.; Zava, O.; Süss-Fink, G.; Juillerat-Jeanneret, L.; Therrien, B. *J. Biol. Inorg. Chem.* **2009**, 14, 101-109.

4) Pernot, M.; Bastogne, T.; Barry, N. P. E.; Therrien, B.; Koellensperger, G.; Hann, S.; Reshetov, V.; Barberi-Heyob, M. *J. Photochem. Photobiol. B* **2012**, 117, 80-89.

5) Naik, A.; Rubbiani, R.; Gasser, G.; Spingler, B. *Angew. Chemie Int. Ed.* **2014**, 53, 6938-6941.

RNA internal loop: suitable binding site for metallo-intercalators?E. Alberti¹, M. Coogan², D. Donghi^{1*}¹University of Zurich, ²Lancaster University

Metal complexes are widely used for therapeutic and diagnostic applications, and their interaction with DNA has been extensively studied [1,2]. However, besides DNA, there are several other possible biological targets, including RNA, which is particularly attractive owing to its involvement in several biological processes and its complex structural diversity [3]. Our project deals with the study of the RNA interaction of a family of Re(I)dppz complexes with potential as bio-imaging agents, whose DNA binding properties were already studied [4]. It was also shown that their cellular uptake varies upon changing their axial ligand and that they can accumulate in DNA and RNA rich regions [4]. However, there are no information on their RNA interaction. Therefore we are investigating the behaviour of one of these complexes, namely $[\text{Re}(\text{CO})_3(\text{dppz})(3\text{-CH}_2\text{OH-Py})]^+$, in the presence of two different RNA constructs containing common RNA secondary structural features. The first construct is a 27 nucleotide long RNA, derived from the mitochondrial group II intron ribozyme Sc.ai5y [5], which contains a GU wobble, an internal and a terminal loop. The second one is an analogous short RNA that does not contain the internal loop and is used to probe a preferential interaction for this feature. A similar DNA sequence has also been used in the binding assays. Moreover, the behaviour of a classical ruthenium complex, known to act as DNA molecular "light switch" [6], was also studied for the sake of comparison. The RNA/DNA binding was studied by UV/Vis and fluorescence titrations, as well as ethidium bromide fluorescence displacement assays. Also, NMR experiments were performed to localize the interaction site. Interestingly, our NMR data suggest, in addition to an overall generalized binding, a localized interaction at the RNA internal loop. Further experiments are currently being performed to better understand the nature of this interaction.

Financial support by the Swiss National Science Foundation (Ambizione fellowship PZ00P2_136726 to DD), by the University of Zurich (including the Forschungskredit grant FK-13-107 to DD) and within the COST Action CM1105 is gratefully acknowledged.

[1] Michael P. Coogan, Vanesa Fernández-Moreira, *Chemical Communications*, **2014**, 50, 384-399.

[2] Dik-Lung Ma, Hong-Zhang He, Ka-Ho Leung, Daniel Shiu-Hin Chan, Chung-Hang Leung, *Angewandte Chemie International Edition*, **2013**, 52, 7666-7682.

[3] Lirui Guan, Matthew D. Disney, *ACS Chemical Biology*, **2012**, 7, 73-86.

[4] Flora L. Thorp-Greenwood, Michael P. Coogan, Lallan Mishra, Niraj Kumari, Greta Rai, Srikrishna Saripella, *New Journal of Chemistry*, **2012**, 36, 64-72.

[5] Daniela Donghi, Maria Pechlaner, Cinzia Finazzo, Bernd Knobloch, Roland K.O. Sigel, *Nucleic Acids Research*, **2013**, 41, 2489-2505.

[6] Alan E. Friedman, Jean Claude Chambron, Jean Pierre Sauvage, Nicholas J. Turro, Jacqueline K. Barton, *Journal of American Chemical Society*, **1990**, 112, 4960-4962

Secrets of *in vitro* RNA folding and splicing revealed by fluorescent PNA labels

Ilija Vukadin¹, S. Zelger-Paulus¹, M. Gulotti-Georgieva¹, A. G. Schmitz¹, R. K. O. Sigel^{1*}

¹University of Zurich

RNA is a versatile and functionally highly diverse molecule involved in numerous crucial processes within biological systems. It's mostly known for its catalytic activity and role as carrier of genetic information. The latter comprises the maturation of RNA, of which RNA splicing is an essential process not yet completely understood.

A model system of high interest is the wild type group II intron ai5 from *Saccharomyces cerevisiae*. The self-catalytic property of this ribozyme can *in vitro* be induced by a high non-physiological concentration of Mg²⁺ ions, whereas *in vivo* it requires the help of chaperone proteins [1]. Visualization of the *in vitro* splicing event under physiological conditions was enabled by native Polyacrylamide Gel Electrophoresis (PAGE) and Förster Resonance Energy Transfer (FRET) experiments using fluorescent Peptide Nucleic Acid (PNA) labels [2]. PNAs are synthetic molecule with high affinity to nucleic acids and a low mismatch tolerance in complementary PNA/nucleic acid duplexes [3]. Based on these *in vitro* experiments, we want to go further towards *in vivo* studies of the splicing process and improve the comparability of *in vitro* and *in vivo* experiments on RNA macromolecules.

Financial support from the University of Zurich (to RKOS and Forschungskredit to SZP) and the ERC (to RKOS) is gratefully acknowledged.

[1] Anna M. Pyle, Olga Fedorova, Christina Waldsich, *Elsevier*, **2007**, 32 (3), 138 - 145.

[2] Anita G. Schmitz, Susann Zelger-Paulus, Gilles Gasser, Roland K.O. Sigel, *ChemBioChem*, **2015**, DOI: 10.1002/cbic.201500180.

[3] Peter E. Nielsen, Michael Egholm, *Curr Issues Mol Biol*, **1999**, 1(2), 89 - 104.

RNA and Oxaliplatin: Investigation of Possible Platinum Binding SitesM. Zampakou¹, D. Donghi^{1*}¹University of Zurich

Nucleic acids are important intracellular targets for many drugs. For example, platinum(II) anticancer drugs are known to primarily exert their activity upon covalent binding to DNA purine nitrogens [1]. In this context, to better understand their mechanism of action, several studies have been dedicated to the characterization of platinum-DNA adducts [2]. However, also RNA is an important biomolecule, with many important functions *in vivo*. The disruption of these processes can have serious consequences, and it has been already reported that some RNA related activities can be inhibited upon platinum drug administration. Still, only little is known on the effect of platinum drugs on RNA structure and biology [3, 4].

In order to understand how platinum drugs affect RNA structure, we are studying the interaction of a 27 nucleotide long RNA construct, derived from the mitochondrial group II intron ribozyme Sc.ai5γ [5] and used as model system, with oxaliplatin. The latter is an FDA approved anticancer drug used in clinics worldwide [6]. After having found the optimal conditions to obtain and isolate monoplatinated RNA samples, we are now investigating which are the possible platinum binding sites, using a combination of NMR studies as well as hydrolytic and enzymatic digestions of ³²P 5'- and 3'-end labelled samples.

Financial support by the Swiss National Science Foundation (Ambizione fellowship PZ00P2_136726 to DD), by the University of Zurich and within the COST Action CM1105 is gratefully acknowledged

1. Rebecca A. Alderden, Matthew D. Hall, Trevor W. Hambley, J. Chem. Educ., **2006**, 83, 728-734.
2. Jan Reedijk, Proc. Natl. Acad. Sci. U. S. A., **2003**, 100, 3611-3616.
3. Erich G. Chapman, Alethia A. Hostetter, Maire F. Osborn, Amanda L. Miller, Victoria J. DeRose, Met. Ions Life Sci., **2011**, 9, 347-377.
4. Alethia A. Hostetter, Maire F. Osborn, Victoria J. DeRose, ACS Chem. Biol., **2012**, 7, 218-225.
5. Daniela Donghi, Maria Pechlaner, Cinzia Finazzo, Bernd Knobloch, Roland K.O. Sigel, Nucleic Acids Res., **2013**, 41, 2489-2504.
6. Amna Ibrahim, Steven Hirschfeld, Martin H. Cohen, Donna J. Griebel, Grant A. Williams, Richard Pazdur, The Oncologist, **2004**, 9, 8-12.

Studies on the mode of action of cationic β -hairpin antibioticsM. Urfer¹, J. A. Robinson^{1*}¹University of Zurich

With increasing bacterial resistance to existing antimicrobial drugs, the need for new antibiotics active against drug-resistant microorganisms is becoming a pressing issue. Conformationally restrained peptidomimetics based on naturally occurring antimicrobial peptides of the innate immune system are gaining in importance, and are interesting starting points in the discovery of new biologically active molecules. Our research focuses on β -hairpins, a recurrent motive that participates in many important biological processes. Several antimicrobial peptides from the innate immune system have been identified having this structural motif, making the synthesis and use of stable structurally constrained peptides that mimic these of great interest. The peptidomimetics contain 12 residues linked to a hairpin-stabilizing D-Pro-L-Pro template, which can be synthesized by solid phase peptide synthesis. Earlier efforts to discover new cyclic cationic antimicrobial peptides that adopt stable β -hairpin structures led to the discovery of L27-11, which has a potent and selective antimicrobial activity against *Pseudomonas* spp. Following the synthesis and screening of libraries of β -hairpin peptidomimetics, several other candidates with good antimicrobial properties and low toxicity were also identified.

As in the case of L27-11, which was shown to target the outer membrane protein LptD and to inhibit its key role in outer membrane biogenesis, understanding the mechanisms of action of these new antimicrobial peptidomimetics is a priority, to identify new targets that could be used to fight against bacterial resistance. In this work, we describe one novel peptidomimetic antibiotic, showing potent activity against *E. coli*. Combining several known techniques to methodically analyse the potential targets in this bacterium, we were able to obtain evidence for a likely mechanism of action. The influence on membrane integrity was examined by following the uptake of Sytox and release from cells of β -lactamase and β -galactosidase. Radiolabelled precursors were used to assess effects on the biosynthesis of macromolecular components of the cell (proteins, RNA, DNA and cell envelope). Various microscopy techniques (transmission and scanning EM and optical STED), photoaffinity labelling and proteomic studies were used to highlight possible targets within the cell. Although his general approach was applied to one antimicrobial peptidomimetic, because of its potent antimicrobial action against Gram-negative bacteria, it could also be employed to systematically analyse targets of other new antimicrobial peptidomimetic compounds.

Robinson, J. A. (2013). Max Bergmann lecture Protein epitope mimetics in the age of structural vaccinology. *Journal of Peptide Science*, 19(3), 127-140.

Srinivas, N., Jetter, P., Ueberbacher, B. J., Werneburg, M., Zerbe, K., Steinmann, J., ... Robinson, J. A. (2010). Peptidomimetic antibiotics target outer-membrane biogenesis in *Pseudomonas aeruginosa*. *Science*, 327(5968), 1010-1013.

Towards *in vivo* splicing of group II intron ai5y

M. Gulotti-Georgieva¹, S. Zelger-Paulus¹, R. K. O. Sigel^{1*}

¹University of Zurich

The group II intron family represents a large class of non-coding RNA elements found in bacteria and organellar genomes of fungi, plants, and lower eukaryotes. Group II introns are known to undergo self-splicing during mRNA maturation. This occurs in a two-step mechanism, where catalysis of successive trans-esterification reaction leads to autocatalytic excision of the intron and concomitantly to exon ligation. Both steps are reversible rendering group II introns mobile genetic elements [1]. *In vivo* the splicing process is supported by cofactors. Our research focuses on the autocatalytic group II intron ai5y from *Saccharomyces cerevisiae*. For its folding and splicing *in vivo* it requires the assistance of the DEAD-box protein Mss116, an ATP dependent helicase [2]. Intensive studies over the past decades have concentrated on a truncated (modified) intron model, whereas we are interested in the wild type molecule. Both *in vitro* and *in vivo* splicing process of the full length construct will be investigated in the presence and absence of Mss116. Two well-known methods will be applied: native Polyacrylamide Gel Electrophoresis (PAGE) and single-molecule Förster Resonance Energy Transfer (smFRET). The fluorescence labeling of the target molecule on specific positions will be performed either using DNA oligonucleotides or alternatively, Peptide Nucleic Acids (PNAs), both carrying fluorescence dyes. Our results will lead to a detailed understanding of the wild type group II intron ai5y folding and splicing pathway.

Financial support by the ERC and the University of Zurich is gratefully acknowledged.

[1] Alan M. Lambowitz and Steven Zimmerly, *Cold Spring Harb Perspect Biol.*, **2011**, 3(8), a0036160.

[2] Olivier Cordin, Josette Banroques, N. Kyle Tanner, Patrick Linder, *Gene*, **2006**, 367, 17-37.

Characterization of group II intronsretrohoming site at the single molecule level.M. Khier¹, D. Kowerko¹, R. K. O. Sigel^{1*}¹University of Zurich

Group II introns are catalytically active RNA molecules: "*Ribozymes*". They cleave single RNA or DNA strand via two transesterification reactions. Group II introns are strictly dependent on Mg^{2+} for proper folding and catalysis [1]. In this study, we used smFRET to characterize the interaction between d3'EBS1* (Exon Binding Site 1) hairpin and its cognate IBS1* (Intron Binding Site 1). The cognate was either an RNA (IBS1*) or DNA (dIBS1*) fragment [2]. These two interactions, i.e. d3'EBS1*-IBS1* and d3'EBS1*-dIBS1*, mimic the recognition interaction required for splicing and for *retrohoming* of group II introns, respectively. NMR studies of these two constructs revealed three binding pockets for M^{2+} [3-4]. We combine structural information from NMR with kinetic and structural data from smFRET to compare our findings of the d3'EBS1*-dIBS1* construct with those of the d3'EBS1*-IBS1* [5].

Our results indicate that the interaction of the d3'EBS1* hairpin with both cognates displays a specific requirement for Mg^{2+} ions at physiological concentration. However, unlike the d3'EBS1*-IBS1* interaction, d3'EBS1*-dIBS1* requires Mg^{2+} ions to promote docking of dIBS1* to d3'EBS1*. In addition, our results show that the two interactions differ slightly in their conformation of the bound state, in a qualitative agreement with the NMR results performed on the same structures [3-4]. The kinetic analysis revealed that the presence of Mg^{2+} leads to a pronounced heterogeneity in the kinetic of both interactions. We corroborate the finding that heterogeneity in the case of by d3'EBS1*-IBS1* results from an incomplete occupancy of M^{2+} binding pockets by Mg^{2+} and Ca^{2+} ions [5] and we propose that kinetic heterogeneity of d3'EBS1*-dIBS1* interaction results as well from an incomplete occupancy of M^{2+} binding pockets by Mg^{2+} ions.

[1]: Roland K O Sigel, *European Journal of Inorganic Chemistry*,**2003**,2281-2292.

[2]: Linhui Julie Su, Peter Z Qin, William J Michels, Anna Marie Pyle, *Journal of Molecular Biology*,**2001**,306, 655-68.

[3]: Daniela Kruschel, Miriam Skilandat, Roland K O Sigel, *RNA*,**2014**,20, 295-307.

[4]: Miriam Skilandat, Roland K O Sigel, *The Journal of Biological Chemistry*,**2014**,30, 20650-63.

[5]: Danny Kowerko, Sebastian L B König, Miriam Skilandat, Daniela Kruschel, Mélodie C A S Hadzic, Lucia Cardo, Roland K O Sigel, *Proceedings of the National Academy of Sciences of the United States of America*,**2015**,112, 3403-08.

Correlation between structure and antimicrobial activity of chitosan-alkyl thiomers - a biological study

S. Conti¹, M. Croce¹, C. Maake¹, G. R. Patzke^{1*}

¹University of Zurich

Chitosan is a natural polymer consisting of N-acetyl-D-glucosamine and β -1,4-linked-D-glucosamine units. Its intrinsic antimicrobial properties lead this compound to be a promising material for many clinical applications.¹ In our previous work we discovered that functionalization of chitosan with thioglycolic acid produces an excellent antimicrobial agent.² In this study, the relation between structure and antimicrobial activity of different chitosan-thiomers has been investigated, particularly we explored the role of the alkyl-chain length of the thio-substituent.

Chitosans with different degrees of deacetylation (87.5-94.2 %) and molecular weights (25-250 KDa) have been used as starting material for the synthesis of chitosan-thioglycolic acid, -thiolactic acid, -mercaptopropionic acid, -2-methyl-3-sulfanylpropanoic acid, -mercaptobutyric acid, -mercaptohexanoic acid and -8-mercaptooctanoic acid. After the chemical characterization of all the chitosan-thiomers their antimicrobial activity was screened against four different bacterial strains, i.e. two Gram negative (*Escherichia coli* and *Pseudomonas aeruginosa*) and two Gram positive (*Streptococcus sobrinus* and *Streptococcus mutans*) with a reliable and optimized MTT viability assay. Further microbiological tests, including minimal inhibitory and bactericidal concentration determination, disk diffusion test and plate counting were performed.

Our study showed that Gram positive strains were more affected by the treatment with the chitosan-thiomers compared to the Gram negative ones. On the other hand, Gram negative bacteria were more sensitive to chitosan-thioglycolic acid and chitosan-mercaptopropionic acid compared to the other synthesized compounds.

This study emphasizes that the functionalization of chitosan, together with the molecular weight and the degree of the deacetylation, may affect the antimicrobial activity. It is clear that a meticulous characterization is a required step for correlating the structure and the antimicrobial activity, especially for polymeric compounds.

[1] Marguerite Rinaudo, *Progress in Polymer Science*, **2006**, 31 (7), 603-632.


[2] Georg Geisberger, Emina B. Gyenge, Doris Hinger, Andres Käch, Caroline Maake, Greta R. Patzke, *Biomacromolecules*, **2013**, 14(4), 1010-1017.

Peptide Shuttle System to Deliver PNAs to their Place of Action

S. Zelger-Paulus, Ilija Vukadin¹, A. Schmitz¹, P. Anstätt¹, G. Gasser¹, R. K. O. Sigel^{1*}

¹University of Zurich

We are exploring a specific process during RNA maturation called splicing and trying to understand the folding mechanism of the involved RNA under native conditions. Thereby, we are applying a new labelling strategy for RNA by using hybridization probes with an unnatural backbone [1]. Peptide nucleic acids (PNAs) are an excellent alternative to DNA oligonucleotides, since they are highly specific and affine to RNA and less prone to enzymatic degradation. Towards *in vivo* splicing studies, it is essential to find an ideal shuttle system to deliver the fluorescent PNA probes to their final intracellular destination, namely to the mitochondrial matrix of *S. cerevisiae*. Hence two lipid bilayers have to be overcome, the cell membrane and the mitochondrial membrane (Figure). RNA complementary PNAs can be designed as follows: i) an N-terminal signal peptide to facilitate a directed transport into the mitochondria [2] and ii) a C-terminal fluorophore to visualize the RNA. These labels have to be actively imported through the outer (TOM) and the inner membrane channels (TIM23) of the mitochondria. After successful entry, the signal-peptide is proteolytically cleaved. We aim to use this strategy to introduce our fluorescently labelled PNA into mitochondria, to specifically target the groupII intron and to investigate the splicing mechanism by FRET. Here we are presenting the first step towards a deeper comprehension of RNA-driven processes occurring in living cells.



Financial support by the ERC, the University of Zurich and the Forschungskredit of the University of Zurich (S. Zelger-Paulus), grant no [FK-13-095] is gratefully acknowledged.

[1] Anita G. Schmitz, Susann Zelger-Paulus, Gilles Gasser, Roland K.O. Sigel, *ChemBioChem*, **2015**, DOI:10.1002/cbic.201500180.

[2] François Sieber, Antonio Placido, Samira El Farouk-Ameqrane, Anne-Marie Duchêne, Laurence Maréchal-Drouard, *Nucleic Acids Res*, **2011**, 39, 14.

Roles of the continuous internal water pathway in G-protein-coupled receptors activations

S. Yuan^{1,2}, H. Vogel^{2*}, S. Filipek^{3*}

¹Actelion Pharmaceuticals Ltd, ²EPF Lausanne, ³The University of Warsaw

More than 800 human GPCRs allow the selective detection of extracellular signals as diverse as photons, odorants, flavors, nucleotides, hormones, neurotransmitters – revealing GPCRs fundamental role in signal transduction. As they regulate many central physiological processes and are thus implicated in many diseases, GPCRs are among the most important targets for modern medicines. In spite of this medical importance and the recent progress in elucidating the 3D structures of various GPCRs, central questions how these receptors recognize extracellular chemical signals and transfer them across the cellular membrane to finally evoke an intracellular response are largely unresolved at a molecular level, mainly because the different steps during signal transmission are not directly accessible by experiments. In this context we are concentrating on central questions of GPCR mediated cellular signalling using molecular dynamics simulations and structural modelling. Our work revealed for the first time, in atomic detail, the entire process of transmembrane signalling of various GPCRs: we found that ligand binding induces a series of conformational changes within a GPCR which opens a gate inside the receptor for water molecules entering the internal region of the receptor and subsequently driving conformational switches within the receptor which finally led to the activation of a G protein on the intracellular side of the receptor. The principle of uncovered internal water channels has successfully applied to the design sub-nM active compounds for two GPCRs targets.


1. Yuan S.*; Palczewski K.; Peng Q.; Kolinski M.; Vogel H.*; Filipek S.* (2015) Angew. Chem. Int. Edt. DOI: 10.1002/anie.201501742R2
2. Yuan S.*; Hu Z.; Filipek S.; Vogel H.* (2015) Angew. Chem. Int. Edt 127(2),566–569
3. Yuan S.*; Filipek S.; Palczewski K.; Vogel H.* (2014) Nature Communication 5, 4377 DOI:10.1038/ncomms5733
4. Yuan S.*; Wu R.; Latek D.; Trzaskowski B.; Filipek S.* (2013) PLoS Computational Biology DOI: 10.1371/journal.pcbi.1003261
5. Yuan S.*; Vogel H.; Filipek S.* (2013) Angew. Chem. Int. Edt 52(38) 10112–10115

Molecular Mechanism of Ruthenium and Gold Anticancer Agents in the Allosteric Regulation of the Nucleosome Core Particle (NCP)

G. Palermo¹, T. Riedel², C. A. Davey³, P. Dyson¹, U. Röthlisberger^{1*}

¹EPF Lausanne, ²University of Zurich, ³Nanyang Technological University, 60 Nanyang Drive, Singapore 637551, Singapore

Metal-based antitumor drugs have emerged as effective chemotherapeutic agents.^[1,2] In particular, Ru-based RAPTA (Ru(II), Arene, PTA = 1,3,5-triaza-7-phosphatricyclo[3.3.1.1]decane) compounds appear to interfere with transcription, resulting in apoptosis.^[3] Besides, the gold-based compound Auranofin – a clinically approved antiarthritic drug – has shown significant antiproliferative properties in in vivo and in vitro models.^[4] Here, we report the molecular mechanism of Auranofin and Rapta-T (T = Toluene) in the allosteric regulation of histone proteins of nucleosome core particles (NCP), the basic repeating unit of chromatin. High-resolution crystal structures indicate that Auranofin and Rapta-T bind at two sites of the NCP histone core that are ~35 Å apart. In spite of this, biochemical and quantitative ICP/MS experiments show a ~3 fold increased binding of Auranofin when in combination with RAPTA-T, therefore suggesting a synergistic effect of the two compounds. Integrating these experimental results with microsecond molecular dynamics (MD) simulations, we are able to propose at the atomistic level the cooperative binding mechanism of Auranofin and Rapta-T. Our results show that the allosteric communication between the ruthenium and gold sites occurs through a series of subtle conformational changes within the protein framework, which are transmitted via coupled motions of the histone components. These findings depict a novel mechanism for the allosteric regulation of the NCP. Overall, our study provides novel insights on the mechanism of action of Ru(II) and Au(I) anticancer agents, potentially leading to the discovery of new therapeutic strategies against cancer.



[1] Gianni Sava, Alberta Bergamo, Paul J. Dyson, *Dalton Transactions*, **2011**, 40, 9069-9075.

[2] Christian G. Hartinger and Paul J. Dyson, *Chem Soc Rev* **2009**, 38, 391-401.

[3] Zenita Adhireksan, Gabriela E. Davey, Pablo Campomanes, Michael Groessl, Catherine M. Clavel, Haojie Yu, Alexey A. Nazarov, Charmian Hui Fang Yeo, Wee Han Ang, Peter Dröge, Ursula Röthlisberger, Paul J. Dyson and Curt A. Davey *Nature Communications*, **2014**, 5, 3462-3474.

[4] Elena Vergara, Angela Casini, Francesca Sorrentino, Olivier Zava, Elena Cerrada, Maria Pia Rigobello, Alberto Bindoli, Mariano Laguna and Paul J. Dyson *ChemMedChem* **2010**, 5, 96-102.

Long-Lived States of Pairs of Fluorine-19 Nuclei: a new Tool for Ligand-Protein Screening

R. Buratto¹, D. Mammoli¹, A. Bornet¹, E. Canet^{1,2}, B. Vuichoud¹, J. Milani¹, S. Jannin¹, G. Bodenhausen^{1,2*}

¹EPF Lausanne, ²ENS Paris

NMR is routinely used in pharmaceutical industry for screening drug candidates by ranking their affinities for protein targets. We have exploited the enhanced contrast afforded by Long-Lived States (LLS) for drug screening [1]. Because T_{LLS} is greater than T_1 , the contrast between the signal intensities of free and bound ligands is dramatically improved, which allows one to reduce the concentrations of both ligands and protein targets. LLS can be used to screen and determine the dissociation constants K_D of molecular fragments that bind weakly to protein targets. We used our LLS method for screening weakly binding fragments against Heat Shock Protein Hsp90, a target for cancer treatment. By exploiting the LLS behavior of a spy molecule, we demonstrated that it is possible to measure dissociation constants K_D as large as 12 mM [2]. This corresponds to a very weak binding regime, where most other biophysical techniques fail, including other NMR methods based on the observation of ligands. This weak binding regime is crucial for fragment-based binding studies (FBB).

More recently, we have combined LLS for drug screening with ¹H dissolution-DNP to enhance the sensitivity. This allows one to reduce the concentrations of ligands and the protein trypsin to 120 μ M and 1.4 μ M respectively [3]. We observed dramatic differences between the spectra of ligands in the presence or absence of trypsin. The presence of a stronger ligand displaces a weaker "spy" ligand, which allows one to determine the dissociation constants K_D of the strong ligand by monitoring the signals of the spy ligand. We demonstrated that it is possible to perform experiments with a very dilute 12 μ M solutions of ligands, obtaining similar signal-to-noise ratios in one-shot DNP experiments as after several hours of acquisition using conventional experiments without DNP.


We are now exploring LLS involving pairs of ¹⁹F nuclei to study binding phenomena. The ¹⁹F spectra do not suffer from any overlapping signals. In a custom-designed fluorinated ligand that binds trypsin, we have observed T_{LLS} = 2.6 s and T_1 = 0.6 s, hence a promising ratio T_{LLS}/T_1 is greater than 4. We found a dramatic effect on the LLS lifetime T_{LLS} of 800 μ M of the fluorinated ligand: an 85% contrast has been observed between signals derived from T_{LLS} with/without 2 μ M trypsin. The titration of the fluorinated ligand against trypsin gave a dissociation constant K_D = 106 μ M. This fluorinated ligand can be used as spy molecule in competition experiments, which allows one to rank the affinities of arbitrary ligands that do not contain any fluorine.

Comparison of ^{18}F -labeled alpha and gamma-conjugated folate derivatives for tumor imaging using positron emission tomography (PET)

S. D. Boss¹, T. Betzel¹, C. Müller², S. Haller², V. Groehn³, C. R. Fischer¹, R. Schibli², S. M. Ametamey^{1*}

¹ETH Zurich, ²Paul Scherrer Institute, Villigen, ³Merck & Cie, Schaffhausen

The folate receptor (FR) is upregulated in various epithelial cancer types (FR α -isoform), whereas healthy tissues show only restricted expression of FR- α . FR-targeted imaging is therefore a promising approach for the detection of FR-positive cancer tissue. Folic acid has two carboxylic functionalities at the alpha (α) and gamma (γ) positions which are accessible for chemical modification. We present the (radio)syntheses, biological evaluation and comparison of three pairs of fluorine-18 labeled α - and γ -conjugated folate derivatives.



Click chemistry was used for the syntheses of the three different pairs of folate regioisomers (α/γ -FDG-, α/γ -fluoroethyl- and α/γ -fluorobutyl-folates). All synthesized folate regioisomers exhibited nanomolar binding affinities to the FR (IC_{50} values in the range of 1 to 3nM) which were determined using FR-expressing KB tumor cells. PET and biodistribution studies revealed similar KB-tumor uptake for both the α -regioisomers and their corresponding γ -isomers of all three pairs of folate-conjugates (α -[^{18}F]**1**: $10.9 \pm 0.52\%$ ID/g, γ -[^{18}F]**1**: $9.05 \pm 2.12\%$ ID/g; α -[^{18}F]**2**: $12.5 \pm 1.04\%$ ID/g, γ -[^{18}F]**2**: $7.24 \pm 0.99\%$ ID/g; α -[^{18}F]**3**: $3.73 \pm 0.62\%$ ID/g, γ -[^{18}F]**3**: $3.22 \pm 1.05\%$ ID/g). In contrast, liver uptake was more than two times higher for the γ -conjugated derivatives compared to their corresponding α -derivatives (α -[^{18}F]**1**: $3.01 \pm 0.48\%$ ID/g, γ -[^{18}F]**1**: $8.37 \pm 1.19\%$ ID/g; α -[^{18}F]**2**: $2.77 \pm 0.57\%$ ID/g, γ -[^{18}F]**2**: $8.66 \pm 1.19\%$ ID/g; α -[^{18}F]**3**: $0.90 \pm 0.22\%$ ID/g, γ -[^{18}F]**3**: $2.25 \pm 0.15\%$ ID/g). A significant difference between the α - and γ -regioisomers was also observed for the kidney uptake values, whereby the γ -regioisomers showed a 50% lower uptake compared to their α -analogues. These results clearly demonstrated that the site of conjugation on the glutamyl moiety of folic acid has a significant impact on the *in vivo* behavior of radiofolates but not on their *in vitro* FR-binding affinity.

- [1] S. D. Weitman et al, *Cancer Res.* **1992**, 52, 6708-6711.
- [2] C. Müller, *molecules*, **2013**, 18, 5005-5031.
- [3] T. Ross et al, *Bioconjugate Chem.* **2008**, 19, 2462-2470.
- [4] C. R. Fischer et al, *Bioconjugate Chem.* **2012**, 23, 805-813.

Human Biomonitoring Bridging Gaps between Medicine and EnvironmentB. Shomar¹

¹Qatar Environment and Energy Research Institute (QEERI), Qatar Foundation, POBox 5825,
Doha, Qatar

Keywords: environmental quality, human health, human biomonitoring.

Bringing the two schools of clinical medicine and environmental sciences is always challenging. Elevated and escalated diseases related to the deterioration of environmental quality call for some research areas in between. These areas can address the roots of the environmental problems (identification of contaminants, natural and anthropogenic contribution, new sensitive and selective analytical tools, contaminant Chemodynamics, transfer to the food chain, etc.) and occurrence in the human body showing short or long term symptoms up to the gene or molecular levels. Human biomonitoring (HBM) is a very powerful tool on exposure sciences. Briefly, HBM is defined as the measurement of concentrations of chemicals or their metabolites in human biological samples such as blood, serum, urine, saliva, breast milk, nails, hair or body parts and tissues. It includes Environmental Health Surveillance and Environmental Health Tracking.


The complexity of environment and the emerging contaminants should be associated with dynamic tools for monitoring. Theory (modeling, computation and simulation) is not driven by or linked to the experimental work and coordination is absent or minimum. A very strong roadmap should be known to bridge the area of environmental deterioration and human health. Qualified and certified national laboratories and competent staff are needed to target ultra-trace inorganic and organic contaminants from sources to fate. Contaminant speciation, metabolites, isotopes and derivatives are examples. The presentation covers new meanings of environmental monitoring and highlights risks associated with contaminants.

Palladium-mediated Suzuki-Miyaura coupling: an efficient method for the formation of therapeutically relevant protein conjugatesA. Dumas^{1,2}, P. Couvreur¹, B. G. Davis²¹Université Paris-Sud, ²University of Oxford

Chemical protein modification is a powerful tool for probing natural systems structure and function and generating protein conjugates with new or improved properties.[1] Owing to their broad functional group tolerance and biological compatibility, palladium-catalyzed reactions targeting unnatural amino acids emerged as attractive strategies for site-specific protein modification.[2]

The efficiency and robustness of these reactions was recently demonstrated for the PEGylation of proteins, a widespread approach to improve the stability and pharmacokinetic properties of protein drugs. The development of palladium-mediated *Suzuki-Miyaura* coupling at genetically encoded or chemically installed halogenated amino acids enabled the efficient synthesis of structurally defined protein-PEG conjugates under physiological conditions.[3] Thanks to the liganding properties of the PEGylated boronic acids involved, this biotechnologically relevant reaction proceeds in a “self-ligated” manner, using the simplest palladium salts as catalyst, without the need for exogenous ligands.[3]

Given the biorthogonality and the high specificity of the *Suzuki-Miyaura* reaction as well as the remarkable properties of palladium in biological settings, and its low associated toxicity, exciting possibilities emerge for the application of this reaction to more complex biological systems.[4]



[1] Anaëlle Dumas, Lukas Lercher, Christopher D. Spicer, Benjamin G. Davis, *Chemical Science*, **2015**, 6, 50-69.

[2] Christopher D. Spicer, Benjamin G. Davis, *Nature Communication*, **2014**, 5:4740.

[3] Anaëlle Dumas, Christopher D. Spicer, Zhanghua Gao, Tsuyoshi Takehana, Yuya A. Lin, Tohru Yasukohchi, Benjamin G. Davis, *Angewandte Chemie International Edition*, **2013**, 52, 3916-3921.

[4] Anaëlle Dumas, Patrick Couvreur, *Chemical Science*, **2015**, 6, 2153-2157.

Enzyme catalyzed sulfur-carbon bond formation by ergothioneine biosynthetic sulfoxide synthaseK. Goncharenko¹, F. Seebeck^{1*}¹University of Basel

EgtBs form a class of non-heme iron enzymes, which catalyze oxygen-dependent sulfur-carbon bond formation between low molecular weight thiols and *N*- α -trimethyl histidine as the central step in ergothioneine biosynthesis (Fig. 1). The crystal structure of EgtB from *Mycobacterium thermoresistibile* in complex with gamma-glutamylcysteine and *N*- α -trimethyl histidine implicate both substrates and three histidine residues as ligands in an octahedral iron binding site. In the secondary coordination sphere we identified a tyrosine residue which may serve as a hydrogen-bond or proton donor to an iron (III)-superoxo or -peroxo species. Mutation of this residue to phenylalanine produced a variant with 300-fold reduced sulfoxide synthase activity. Instead, this protein catalyzes thiol dioxygenation with an efficacy that rivals naturally evolved cysteine dioxygenases (CDO). Comparison of the EgtB and the CDO structures reveals two very similar active sites within two entirely unrelated protein folds. EgtB likely forms the same iron (III)-superoxo intermediate as described for CDO, but strategic placement of a tyrosine residue as catalytic acid promotes sulfur-carbon bond formation and prohibits thiol dioxygenation.





Figure 1. Biosynthesis of ergothioneine in *M. thermoresistibile* via sulfoxide intermediate.

Electrostatic Effect Of Halogenation On The Thermodynamic Stability Of Rapid Insulin Analogs

K. El Hage¹, M. A. Weiss², M. Meuwly^{1*}

¹Department of Chemistry, University of Basel, ²Department of Biochemistry, Case Western Reserve University, Cleveland, Ohio

Insulin analogs are designed containing amino acid substitutions at or near the classical dimerization surfaces of the zinc insulin hexamer [1,2]. One of these amino acids is Phe^{B24}, an invariant aromatic anchor at this interface and site of a human mutation causing diabetes mellitus (See figure below [3]). A recent study showed that a nonstandard aliphatic mutagenesis at this position can optimize the molecular properties of therapeutic proteins and lead to accelerated hexamer disassembly while preserving native receptor binding affinity and biological activity [3]. These findings motivated us to investigate further substitutions by halogenated aromatic rings. The effect of these mutations on the conformational ensemble and on the differential solvation free energy is evaluated using thermodynamic integration with the CHARMM force field and results are compared to experimental data. In addition, we used the multipolar module [4] that allows us to study the critical dependence of the halogen-induced electrostatic dipole on the position of the halogenation (ortho, meta or para) and its role in adding solvation free energy changes. Preliminary results show, that in addition to local chain stabilization, a global thermodynamic stabilization of 1 kcal/mole takes place reflecting enhanced enthalpic interactions via weakly polar interactions by the modified aromatic ring and a more favorable solvation free energy.



Since insulin provides a model for the therapeutic application of protein engineering, the resulting differential effects of these modifications can help provide a molecular strategy to augment protein stability while preserving rapid action.

[1] Zoete, V.; Meuwly, M. *J. Comput. Chem.* **2006**, 27, 1843-1857.

[2] Pandeyarajan et al. *J. Bio. Chem.* **2014**, 289, 23367-23381.

[3] Pandeyarajan et al. *J. Bio. Chem.* **2014**, 289, 34709-34727.

[4] Bereau, T.; Kramer, C.; Meuwly, M. *J. Chem. Theory Comput.* **2013**, 9, 5450-5459.b

Preventing aggregation of porphyrinic photosensitizers using a biodegradable triblock copolymer

I. Gjuroski¹, M. Vermathen¹, J. Furrer^{1*}

¹University of Bern

Photodynamic therapy (PDT) is a form of phototherapy using nontoxic light-sensitive compounds that are exposed selectively to light, upon which they become toxic to targeted malignant cells (phototoxicity). Over the last few decades porphyrins have gained much interest as promising photosensitizers but their intrinsic hydrophobicity and tendency to aggregate in aqueous solution represent limiting factors which decrease their PDT efficacy. One approach to overcome these problems is the use of drug delivery vehicles [1]. Among those vehicles, some polymers like polyvinylpyrrolidone (PVP) have received much attention, notably because they are nontoxic as well as soluble in water. It was also shown that the biological activity and phototoxicity of porphyrins can be enhanced by attaching an amino acid to form porphyrin amino acid conjugates [2].

The aim of this study was to probe the propensity of the biodegradable triblock copolymer Kolliphor P188 to disaggregate various chlorin e6 amino acid conjugates (xCE) [3].

The interaction and loading between the copolymer Kolliphor P188 and the xCE conjugates was probed using NMR spectroscopy as a main tool. The results indicate that Kolliphor P188 is able to monomerize all xCE conjugates in aqueous buffer at room temperature. NMR diffusion data suggest that the loading depends on the structure of the amino acid residue.

[1] Bechet D. et al, *Trends Biotechnol.* **2008**, 26, 612.

[2] Jinadasa, R. G. W. et al, *J. Med. Chem.* **2011**, 54, 7464.

[3] Vermathen M. et al, *J. Phys. Chem.* **2013**, 117, 6990–7001.

Why are Vesicles of the Artificial 1,3-Diamidophospholipid Pad-PC-Pad Mechanosensitive?

D. Müller¹, A. Zumbühl^{1*}

¹University of Fribourg

We have shown that vesicles of the artificial phospholipid Pad-PC-Pad (**1**) are lenticularly shaped and mechanosensitive.¹Temperature, ultrasound, enzymes, pH-changes, redox-reactions or photo-reactions are all well known as trigger for cargo release from phospholipid liposomes.²On the other hand mechanosensitivity is a fresh concept in targeted drug delivery. If vesicles of **1** are exposed to increased shear-stress they release their content. The change in shear stress in blocked arteries could be used as a trigger for targeted drug delivery for the acute treatment of a myocardial infarction patient.





Figure 1: left: chemical structure of the 1,3-diamidophospholipid Pad-PC-Pad (**1**), right: cryo transmission electron micrograph of extruded unilamellar Pad-PC-Pad vesicles, the scale bar is 200 nm.

Mechanosensitivity is based on the unnatural 1,3-fatty acyl chain substitution pattern found in **1** which allows a tail interdigitation of the opposing membrane leaflets. The interdigitation increases the lateral stiffness of the bilayer in a vesicle and rapid cooling from temperatures above T_m leads to the formation of lenticularly shaped vesicles. To proof this hypothesis and to gain a deeper understanding of the mechanosensitivity of phospholipid vesicles we synthesized different artificial lipids showing the natural 1,2- or the unnatural 1,3-fatty acyl chain substitution pattern each with and without the ability to form hydrogen bonds. Preliminary results indicate that hydrogen bonding is less important for mechanosensitivity than the 1,3-substitution pattern in Pad-PC-Pad.

[1] M. N. Holme, I. A. Fedotenko, D. Abegg *et al.*, *Nature Nanotechnol.*, **2012**, 7, 536-543.

[2] D. Mellal, A. Zumbuehl, *J Mater. Chem. B*, **2014**, 2, 247-252.

Addressing the tempormandibular joint disorderE. Stalder¹, A. Zumbühl^{1*}¹University of Fribourg

The temporomandibular joints (TMJ) are the joints connecting the lower jaw (mandible) to the skull and regulating the action of the mandible. Temporomandibular joint disorder (TMD) is a generic term designating the dysfunction of the TMJ. TMD is caused by a broad range of causes such as injuries of the jaw or the TMJ, disc displacement, osteoarthritis, reumathoid arthritis, teeth grinding. The current treatment ranges from physical therapy to surgical replacement of the joint, with the typical treatment being the administration of a painkiller with all the known side effects.[\[1\]](#)

The work presented here aims at fulfilling the need for a mid- to long-term alternative to the parental or oral administration of a painkiller. The idea is to design and study an *in situ* crosslinking hydrogel containing drug loaded liposomes for a long-term and on demand drug delivery system. Periodate oxidized dextran is reacted with amine bearing liposomes and polyethylenimine in order to form a hydrogel. The chemistry and biophysical properties of this gel will be presented.

[1] Ingawale, S.; Goswami, *Ann. Biomed. Eng.* 2009, 37, 976-996.


NMR investigation of the human RNA BCL2 G-quadruplex: restricting folding dynamics

A. Dominguez-Martin¹, R. K. O. Sigel^{1*}

¹University of Zurich, Department of Chemistry, Winterthurerstrasse 190, Zurich, CH-8057 Zurich

A 25 nucleotide guanine rich sequence (5'-G₅CCGUG₄UG₃AGCUG₄-3') was found in the 5'-untranslated region of the mRNA of the human BCL2 proto-oncogene (B-Cell Lymphoma/Leukemia-2). This sequence is prone to fold into a specific non-canonical secondary structure, a so-called G-quadruplex (G4). Interestingly, the presence of this G4 structure has shown to down-regulate the expression of the bcl2 protein *in cellulo* [1]. The BCL2 family of proteins play a crucial role in the regulation of apoptosis [2]. For instance an aberrant expression of the bcl2 protein has been isolated in several human cancers. Therefore, the understanding of this biologically relevant RNA G4 sequence is of high interest for anticancer therapy.

1D ¹H-NMR spectra of the 25-nt BCL2 G4-forming sequence in the presence of K(I) ions revealed the formation of multiple G4 structures. To reduce the dynamics of the system, the sequence was shortened to yield a more restricted 22-nt sequence (5'-G₃CCGUG₄UG₃AGCUG₃-3'). Our results showed the stabilization of mainly one G4 structure upon the addition of K(I), defined as unimolecular, intramolecular and parallel-stranded. However, additional NMR data evidenced the presence of two different but non-equally populated species in solution. This fact hampers NMR investigations. Thus, further stabilization of the 22-nt BCL2 construct was performed by mutational studies. The aim is to isolate one single G4 conformation, which allows us to determine the solution structure of this G-quadruplex by NMR. The triple mutated sequences BCL2_6A8U17U and BCL2_6A8A17U successfully stabilize one single G-quadruplex structure with different loop isomers. Preliminary NMR data shows promising results regarding the structural elucidation of the BCL2 G-quadruplex.



Financial support by the Fundacion Ramon Areces and Forschungskredit (A.D.M.), the Swiss National Science Foundation (R.K.O.S.), the COST Action CM1105 (Swiss State Secr. Edu. Res. Innov. to RKOS) and by the University of Zurich is gratefully acknowledged.

- [1] Anthony Bugaut, Shankar Balasubramanian, *Nucleic Acids Research*, **2012**, 40, 4727-4741.
[2] Jerry M. Adams, Suzanne Cory, *Science*, **1998**, 28, 1322-1326.

Metal ion dependency and multimerization behavior of biologically relevant human RNA G-quadruplexes


H. Guiset-Miserachs¹, R. Börner¹, D. Donghi¹, S. Johannsen¹, R. K. O. Sigel^{1*}

¹Department of Chemistry, University of Zurich, Winterthurerstrasse 190, 8057 Zurich

Guanine-rich nucleic acids fold into non-canonical helical structures, called G-quadruplexes (G4), and formed by stacking of guanine planes. RNA G4s appear in regulatory regions of the transcriptome, suggesting they play a role *in vivo*, and indeed their presence in human cells was proven [1]. We study a sequence from the 5' UTR of the NRAS oncogene mRNA (5'-G₃AG₄CG₃UCUG₃-3'), shown to inhibit translation *in vitro* and to be accessible for antitumor targeting [2, 3].

G4 motifs need charge-balancing metal ions to fold and remain stable. We investigated the influence of the metal ion type and concentration on G4 folding by following two G4 sequences with several mono- and divalent metal ions (see figure): the NRAS and the telomeric-repeat containing RNA (TERRA, 5'-(UUAG₃)₄), the most studied G4 RNA so far [4].

Circular dichroism showed that Na⁺, K⁺, Sr²⁺ and Ba²⁺ promote G4 folding, while only K⁺ and Sr²⁺ yield further stabilized G4s, as seen by a significant increase in melting temperatures. Further addition of K⁺ and Sr²⁺ leads to unspecific interactions and cation-mediated multimers, evidenced by polyacrylamide gel electrophoresis, dynamic light scattering and mass spectrometry. Comparison with published DNA G4s shows G4 RNAs to be more stable, more prone to multimerize and less Na⁺-sensitive. These studies delve into the under-studied RNA G4 motifs and their folding behavior, relevant towards possible functions *in vivo*.



Financial support by the COST Action CM1105, an ERC Starting Grant 2010 (R.K.O.S.), the SNSF (R.K.O.S.), a UZH Forschungskredit (FK-13-090 to H.G.M.) and by the UZH is gratefully acknowledged. We are grateful to Mr. A. Marchand, Dr. F. Rosu and Dr. V. Gabelica (Univ. of Bordeaux) for MS measurements, and Dr. B. Klejevskaja and Prof. R. Vilar (Imperial College London) for collaboration regarding PAGE gels.

[1] Giulia Biffi, Marco Di Antonio, David Tannahill, Shankar Balasubramanian, *Nat. Chem.*, **2014**, 6, 75-80.

[2] Sunita Kumari, Anthony Bugaut, *et al.*, *Nat. Chem. Biol.*, **2007**, 3, 218-221.

[3] Anthony Bugaut, Raphaël Rodriguez, Sunita Kumari, *et al.*, *Org. Biomol. Chem.*, **2010**, 8, 2771-2776.

[4] Brian Luke, Joachim Lingner, *EMBO J.*, **2009**, 28, 2503-2510.

Structure - antimicrobial activity relationships of chitosan-alkyl thiomers

M. Croce¹, S. Conti¹, C. Maake¹, G. R. Patzke^{1*}

¹University of Zurich

Chitosan is a biopolymer consisting of N-acetyl-D-glucosamine and β -1,4-linked-D-glucosamine units with wide potential for clinical applications¹ mainly based on its antimicrobial properties. Recently, a new versatile antimicrobial agent has been synthesized by Geisberger et al.², functionalizing chitosan with thioglycolic acid through amide bond formation. In this study, the relationship between the structure and the antimicrobial activity has been explored synthesizing chitosan-alkyl thiomers with different alkyl-chain length.

Chitosans with different degrees of deacetylation (87.5-94.2 %) and molecular weights (25-250 KDa) have been functionalized with thiol-bearing compounds adapting the literature protocols³ to achieve different degrees of substitution. The series encompasses chitosan-thioglycolic acid, -thiolactic acid, -mercaptopropionic acid, -2-methyl-3-sulfanylpropanoic acid, -mercaptobutyric acid, -mercaptohexanoic acid and -8-mercaptooctanoic acid. All the thiomers were characterized by the following techniques: ¹H-NMR, FT-IR, Ellman test, GPC, elemental analysis and ζ -potential measurements. The antimicrobial activity of the title compounds was screened against four different bacterial strains, namely two Gram negative (*Escherichia coli* and *Pseudomonas aeruginosa*) and two Gram positive (*Streptococcus sobrinus* and *Streptococcus mutans*) with a reliable and optimized MTT assay. For the most promising compounds further microbiological studies (minimal inhibitory and bactericidal concentration determination, disk diffusion test and plate counting) were performed.

Chitosan-thiomers showed higher antimicrobial activity against Gram positive strains compared to the Gram negative ones. Besides, Gram negative strains were more affected by chitosan-thioglycolic acid and chitosan-mercaptopropionic acid compared to the other thiomers.

This study gives good evidences that antimicrobial activity may be influenced by the functionalization, together with the molecular weight and the degree of the deacetylation. Moreover, an accurate characterization of the new polymers is required to establish systematic structure-antimicrobial activity relationship.

[1] Chunmeng Shi, Ying Zhu, Xinze Ran, Meng Wang, Yongping Su, Tianmin Cheng, *The Journal of surgical research*, **2006**, 133 (2), 185-192.

[2] Georg Geisberger, Emina B. Gyenge, Doris Hinger, Andres Käch, Caroline Maake, Greta R. Patzke, *Biomacromolecules*, **2013**, 14 (4), 1010-1017.

[3] Constantia E. Kast, Andreas Bernkop-Schnürch, *Biomaterials*, **2001**, 22 (17), 2345-2352.

Real-time Characterisation of a Large, Catalytic and Dynamic RNA by Single Molecule Microscopy.


M. C. Hadzic¹, S. Zelger-Paulus¹, R. K. O. Sigel^{1*}

¹University of Zurich

Group II introns are large natural RNA sequences able to self-excise from the nucleic acid strand via a process called *splicing*, allowing the molecule to fold into a functional conformation. Although tertiary structures and autocatalytic properties of engineered constructs have been unveiled along with bulk and single molecule techniques [1,2], there is little knowledge available about the correlation of splicing events with conformational dynamics, which renders this mechanistic picture incomplete. The present study intends to lift the veil on these folding choreographies and on their role in the reaction efficiency.

The ai5y group IIB intron from the yeast *Saccharomyces cerevisiae* flanked by its original 45 nt 5'-exon was fluorophore-tagged on two sites according to different labelling schemes [3]. Single molecules were trapped into small and surface-immobilized phospholipid vesicles, affording to track the 5' cleavage reaction and its products as well as conformational changes by following the distance-dependant energy transfer (smFRET) from the donor (Cy3) to the acceptor (Cy5) fluorophore.

By means of electrophoresis, the splicing reaction was proven to be inhibited at room temperature under ideal salt concentrations but spontaneously triggered by temperature increase to 40 °C, later extended to sm-FRET investigations. Monoencapsulation and subsequent imaging of large fluorophore-bilabelled systems was successful. Data were evaluated with our recently developed and freely available graphical user interface, allowing complete data evaluation [4]. Our sm-FRET results reveal detailed insights in streaming a biochemical process at the single molecule level, by correlating conformational motion with biological activity.



Funding: ERC Starting Grant (R.K.O. Sigel), UZH Forschungskredit (M.C.A.S Hadzic and S. Zelger-Paulus)

[1] Anna M. Pyle. *Crit. Rev. Biochem. Mol. Biol.* 45 (2010):215-232.

[2] Miriam Steiner, Krishanthi S. Karunatilaka, Roland K.O. Sigel, David Rueda. *PNAS* 105 (2008):1807.

[3] Anita G.Schmitz, Susann Zelger-Paulus, Gilles Gasser, Roland K.O. Sigel. *ChemBioChem* (2015), *accepted*.

[4] Mélodie C.A.S. Hadzic, Danny Kowerko, Richard Börner, Sebastian L.B. König, Mario Heidernätsch, Roland K.O. Sigel. *manuscript in preparation*

Pore size matters - A crowding study of ribozyme folding and activity


R. Börner¹, E. Fiorini¹, R. K. O. Sigel^{1*}

¹University of Zurich

Catalytic RNAs, like the group IIB intron ribozyme of *S. cerevisiae*, are known to require a high magnesium(II) concentration to show folding and function *in vitro* [1]. In contrast, *in vivo* conditions are characterized by a highly crowded cellular environment and much lower ion concentration. Nowadays, molecular crowding agents like poly(ethylene)glycol (PEG), are a widespread tool to mimic cellular crowding and thus, to reach near physiological conditions [2]. However, particular physical and/or chemical properties explaining the crowders influence are mostly not understood. In this study, we gain new insights on how polymer properties like viscosity, pore size etc. influence the activity and folding of a large RNA.

We combined bulk activity assays and single-molecule Förster Resonance Energy Transfer (smFRET) experiments to investigate the influence of a crowding agent on group II intron folding. A screening of PEG volume fraction (%) and molecular weight (MW) was further used to elucidate the impact of different polymer properties on the ribozyme activity.

Our results revealed that upon the influence of a crowding agent, a compaction of the underlying RNA structure is dependent on the volume fraction of the crowding co-solute. Activity assays in the presence of different PEG MW and % unveiled an optimal pore size in terms of catalytic activity, which coincides with the dimension of the folded ribozyme. In summary, an increasing density of the crowding environment shifts the RNA conformations towards the most compact state, but the ribozyme is only active if the crowders network matches its size [4]. We therefore interpret the most compact state as necessary, but not sufficient, to keep the ribozyme active.



Acknowledgement: Financial support from the European Research Council (MIRNA N° 259092, to RKOS), the Swiss National Fund (SNF), and the Forschungskredit Grant of the University of Zürich (FK-14-096 to RB) are gratefully acknowledged.

[1] J.F. Swisher, L.J. Su, M. Brenowitz, V.E. Anderson, A.M. Pyle, *J. Mol. Bio.*, **2002**, 315, 297-310.

[2] D. Kilburn, J.H. Roh, L. Guo, R.M. Briber, S.A. Woodson, *JACS*, **2010**, 132, 8690-6.

[3] M. Steiner, K.S. Karunatilaka, R.K.O. Sigel, D. Rueda, *Proc. Natl. Acad. Sci. U.S.A.*, **2008**, 105, 13853-8.

[4] ^aR. Börner, E. Fiorini, R.K.O. Sigel, *Chimia*, **2015**, 69, 207-212. ^bE. Fiorini, B. Paudel, R. Börner, D. Rueda, R.K.O. Sigel, *to be submitted*.

[5] S.L.B. König, M. Hadzic, E. Fiorini, R. Börner, D. Kowanko, W.U. Blanckenhorn, R.K.O. Sigel, *PLoS ONE*, **2013**, 8, e84157.

The binding mechanism between a B₁₂-specific RNA and its ligand coenzyme B₁₂

S. Gallo¹, A. Musiari¹, P. Choudhary¹, E. Ennifar², F. Steffen¹, P. Dumas², R. K. O. Sigel^{1*}

¹University of Zurich, Winterthurerstrasse 190, 8057 Zürich, Switzerland., ²IBMC-CNRS, Strasbourg

A remarkable characteristic of riboswitches, short RNA elements involved in gene regulation, is their ability to interact directly and specifically with their ligand, mostly being a cellular metabolite. In this context, the interaction between the 202 nt long *btuB* riboswitch with its ligand coenzyme B₁₂ (AdoCbl) is highly interesting due to the complexity of both binding partners.[1, 2]

We present here a comprehensive thermodynamic and kinetic study of the *btuB*-AdoCbl binding mechanism combining extensive isothermal titration calorimetry (ITC) experiments with surface plasmon resonance (SPR) and dynamic light scattering (DLS). The ITC data retrieved from measurements at different conditions were assembled to a global thermodynamic treatment to generate accurate K_D and ΔH -values. Our data propose the existence of two distinct *btuB* species, both binding AdoCbl with high affinity, and their ratio being temperature dependent. This finding was confirmed by SPR. DLS measurements and gel-shift assays conducted at various temperatures complete this study as complementary methods. Based on our data we propose a complex binding mechanism involving a preorganization of the *btuB* riboswitch into two distinct binding-competent forms [3], however similar in their overall, global structure.

[1] A. Nahvi, N. Sudarsan, M. S. Ebert, X. Zou, K. L. Brown, R. R. Breaker, *Chem. Biol.* **2002**, 9, 1043.

[2] S. Gallo, M. Oberhuber, R. K. O. Sigel, B. Kräutler, *ChemBioChem* **2008**, 9, 1408.

[3] P. K. Choudhary, R. K. O. Sigel, *RNA* **2014**, 20, 36.

Financial support by European Research Council (ERC starting grant to R.K.O.S.), the Swiss State Secretariat for Education Research and Innovation (COST Action CM1105), and the University of Zurich is gratefully acknowledged.

Hydrogels with short peptides and their composites with metal nanoparticlesA. Holzheu¹, K. M. Fromm^{1*}¹University of Fribourg, Department of Chemistry, Chemin du Musée 9, 1700 Fribourg

For centuries it is known that silver possesses antimicrobial properties and it was regularly used for the treatment of burns, wounds and several bacterial infections. But with the emergence of antibiotics it was nearly forgotten for almost 50 years^{1,2}.

Nowadays, due to the unique physical and chemical properties of silver nanoparticles and the rising concern regarding infectious diseases induced by multidrug-resistant bacteria, metallic silver has made a remarkable comeback as a potential antimicrobial agent².

Hereby, we propose a silver nanoparticle hydrogel system which can be used for medical purposes against multidrug-resistant bacteria.

Peptide self-assembled hydrogels are a promising class of soft biomaterials for cell culture, regenerative medicine, or drug delivery applications³.

These low molecular weight compounds assemble by weak non-covalent interactions and are responsive to different physicochemical stimuli⁴.

In our case the hydrogel system (see Figure 1) itself can synthesize silver nanoparticles due to its hydrazide end group.

The aim is therefore to synthesize the dipeptide hydrogel Boc(β Ala)₂NHNH₂ and to analyze its chemical, physical and mechanical characteristics with or without the incorporation of silver and to test its suitability for biological applications.




Figure 1: The dipeptide Boc(β Ala)₂NHNH₂

[1] S. Eckhardt, P.S. Brunetto, J. Gagnon, M. Priebe, B. Giese, K. Fromm, *Chem. Rev.*, **2013**, 113 (7), 4708–4754

[2] M. Rai, A. Yadav, A. Gade, *Biotechnology Advances*, **2009**, 27, 76–83

[3] S. Marchesan, L. Waddington, C.D. Easton, D.A. Winkler, L. Goodall, J. Forsythe, P.G. Hartley, *Nanoscale*, **2012**, 4, 6752–6760

[4] M. Tena-Solsona, S. Alonso-de Castro, J.F. Miravet, B. Escuder, *J. Mater. Chem. B*, **2014**, 2, 6192–6197

Micro- and meso-fluidic systems for rapid cell and enzyme characterization

M. P. Marques¹, A. M. Super¹, N. Jaccard¹, N. Dimov¹, Y. Y. Chiang¹, M. J. Davies¹, B. O'Sullivan¹,
N. Szita^{1*}

¹University College London (UCL)

The Bioprocess Microfluidics laboratory of the Department of Biochemical Engineering at UCL (University College of London) is dedicated to leading edge research and development in strategic technological areas focusing on the application of micro fabrication techniques and micro fluidic principles to address challenges and opportunities in the Life Sciences. This includes the system-wide integration and development of analytical methods. The research priorities have special emphasis on Bioprocesses and Regenerative Medicine/Cell Therapy.

Key advantages that make miniaturized systems a cost-effective proposition for early bioprocess development include: significant reduction in reagent use, real-time monitoring and control of culture/reaction variables, ease of sterilization *via* disposable polymer technology, reduced labor due to automation, and the capability to rapidly test different culturing/reaction conditions. Clearly, such micro fabricated systems will be highly significant to address current challenges in bioprocessing and regenerative medicine, in particular if they can accelerate the translation of Life Sciences discoveries to reliable processes for the production of novel medicines.

To achieve this, it is crucial to maintain 'a link' with conventional methods and production systems for the purposes of validation and scale up studies. Further, the devices must be highly versatile to facilitate evaluation and optimization of a wide range of process variables. This versatility must extend beyond the control of the physico-chemical properties of the culture/reaction medium. Depending on the application this may for example include the monitoring (or control) of time profiles of cellular growth to determine growth kinetics, factors secreted by the cells, impact of shear forces, time profiles of product formation (or substrate consumption), the concentration levels of specific metabolites, and others. Ideally, this information is obtained online and in real time to reduce experimental load and to generate truly information-rich experimental data.

In this contribution, we present miniaturised systems for (1) analysis of stem cells for application in regenerative medicine and cellular therapies, (2) continuous culture of microbial cells to find optimal growth conditions for *de novo* engineered cells, and (3) meso-scale enzyme reaction devices to rapidly characterise enzyme systems for asymmetric biocatalysis and production of chiral metabolites.